

Abstract Algebraic Logic – 2nd lesson

Petr Cintula¹ and Carles Noguera²

¹Institute of Computer Science,
Academy of Sciences of the Czech Republic
Prague, Czech Republic

²Institute of Information Theory and Automation,
Academy of Sciences of the Czech Republic
Prague, Czech Republic

www.cs.cas.cz/cintula/AAL

Completeness theorem for classical logic

- Suppose that $T \in \text{Th}(\text{CL})$ and $\varphi \notin T$ ($T \not\vdash_{\text{CL}} \varphi$). We want to show that $T \not\models \varphi$ in some meaningful semantics.
- $T \not\models_{\langle \mathbf{Fm}_{\mathcal{L}}, T \rangle} \varphi$. 1st completeness theorem
- $\langle \alpha, \beta \rangle \in \Omega(T)$ iff $\alpha \leftrightarrow \beta \in T$ (congruence relation on $\mathbf{Fm}_{\mathcal{L}}$ compatible with T : if $\alpha \in T$ and $\langle \alpha, \beta \rangle \in \Omega(T)$, then $\beta \in T$).
- Lindenbaum-Tarski algebra: $\mathbf{Fm}_{\mathcal{L}}/\Omega(T)$ is a Boolean algebra and $T \not\models_{\langle \mathbf{Fm}_{\mathcal{L}}/\Omega(T), T/\Omega(T) \rangle} \varphi$. 2nd completeness theorem
- Lindenbaum Lemma: If $\varphi \notin T$, then there is a maximal consistent $T' \in \text{Th}(\text{CL})$ such that $T \subseteq T'$ and $\varphi \notin T'$.
- $\mathbf{Fm}_{\mathcal{L}}/\Omega(T') \cong \mathbf{2}$ (subdirectly irreducible Boolean algebra) and $T \not\models_{\langle \mathbf{2}, \{1\} \rangle} \varphi$. 3rd completeness theorem

Definition 2.1

Let $\mathbf{A} = \langle A, F \rangle$ be an L-matrix. We define:

- the **matrix preorder** $\leq_{\mathbf{A}}$ of \mathbf{A} as

$$a \leq_{\mathbf{A}} b \quad \text{iff} \quad a \rightarrow^{\mathbf{A}} b \in F$$

- the **Leibniz congruence** $\Omega_{\mathbf{A}}(F)$ of \mathbf{A} as

$$\langle a, b \rangle \in \Omega_{\mathbf{A}}(F) \quad \text{iff} \quad a \leq_{\mathbf{A}} b \text{ and } b \leq_{\mathbf{A}} a.$$

A congruence θ of A is **logical** in a matrix $\langle A, F \rangle$ if for each $a, b \in A$ if $a \in F$ and $\langle a, b \rangle \in \theta$, then $b \in F$.

Theorem 2.2

Let $\mathbf{A} = \langle A, F \rangle$ be an L-matrix. Then:

- 1 $\leq_{\mathbf{A}}$ is a preorder.
- 2 $\Omega_{\mathbf{A}}(F)$ is the largest logical congruence of \mathbf{A} .
- 3 $\langle a, b \rangle \in \Omega_{\mathbf{A}}(F)$ iff for each $\chi \in Fm_{\mathcal{L}}$ and each \mathbf{A} -evaluation e :

$$e[p \rightarrow a](\chi) \in F \quad \text{iff} \quad e[p \rightarrow b](\chi) \in F.$$

Proof.

1. Take \mathbf{A} -evaluation e such that $e(p) = a$, $e(q) = b$, and $e(r) = c$. Recall that in \mathbf{L} we have: $\vdash_{\mathbf{L}} p \rightarrow p$ and $p \rightarrow q, q \rightarrow r \vdash_{\mathbf{L}} p \rightarrow r$. As $\mathbf{A} = \mathbf{MOD}(\mathbf{L})$ we have: $e(p \rightarrow p) \in F$, i.e., $a \leq_{\mathbf{A}} a$ and if $e(p \rightarrow q), e(q \rightarrow r) \in F$, then $e(p \rightarrow r) \in F$ i.e., if $a \leq_{\mathbf{A}} b$ and $b \leq_{\mathbf{A}} c$, then $a \leq_{\mathbf{A}} c$.

Theorem 2.2

Let $\mathbf{A} = \langle A, F \rangle$ be an L-matrix. Then:

- 1 $\leq_{\mathbf{A}}$ is a preorder.
- 2 $\Omega_{\mathbf{A}}(F)$ is the largest logical congruence of \mathbf{A} .
- 3 $\langle a, b \rangle \in \Omega_{\mathbf{A}}(F)$ iff for each $\chi \in Fm_{\mathcal{L}}$ and each \mathbf{A} -evaluation e :

$$e[p \rightarrow a](\chi) \in F \quad \text{iff} \quad e[p \rightarrow b](\chi) \in F.$$

Proof.

2. $\Omega_{\mathbf{A}}(F)$ is obviously an equivalence relation. It is a congruence due to (sCng) and logical due to (MP).

Take a logical congruence θ and $\langle a, b \rangle \in \theta$. Since $\langle a, a \rangle \in \theta$, we have $\langle a \rightarrow^{\mathbf{A}} a, a \rightarrow^{\mathbf{A}} b \rangle \in \theta$. As $a \rightarrow^{\mathbf{A}} a \in F$ and θ is logical we get $a \rightarrow^{\mathbf{A}} b \in F$, i.e., $a \leq_{\mathbf{A}} b$. The proof of $b \leq_{\mathbf{A}} a$ is analogous.

Theorem 2.2

Let $\mathbf{A} = \langle A, F \rangle$ be an L-matrix. Then:

- 1 $\leq_{\mathbf{A}}$ is a preorder.
- 2 $\Omega_{\mathbf{A}}(F)$ is the largest logical congruence of \mathbf{A} .
- 3 $\langle a, b \rangle \in \Omega_{\mathbf{A}}(F)$ iff for each $\chi \in Fm_{\mathcal{L}}$ and each \mathbf{A} -evaluation e :

$$e[p \rightarrow a](\chi) \in F \quad \text{iff} \quad e[p \rightarrow b](\chi) \in F.$$

Proof.

3. One direction is a corollary of Theorem 1.16 and (MP).

The converse one: set $\chi = p \rightarrow q$ and $e(q) = b$: then $a \rightarrow^{\mathbf{A}} b \in F$ iff $b \rightarrow^{\mathbf{A}} b \in F$, thus $a \leq_{\mathbf{A}} b$. The proof of $b \leq_{\mathbf{A}} a$ is analogous (using $e(q) = a$). □

Definition 2.3

An L-matrix $\mathbf{A} = \langle A, F \rangle$ is **reduced**, $\mathbf{A} \in \mathbf{MOD}^*(L)$ in symbols, if $\Omega_A(F)$ is the identity relation Id_A .

An algebra A is **L-algebra**, $A \in \mathbf{ALG}^*(L)$ in symbols, if there is a set $F \subseteq A$ such that $\langle A, F \rangle \in \mathbf{MOD}^*(L)$.

Note that $\Omega_A(A) = A^2$. Thus from $\mathcal{F}i_{\text{Inc}}(A) = \{A\}$ we obtain:

$$A \in \mathbf{ALG}^*(\text{Inc}) \quad \text{iff} \quad A \text{ is a singleton}$$

Examples: classical logic CL and logic BCI

Exercise 3

Classical logic: prove that for any Boolean algebra A :

$$\Omega_A(\{1\}) = \text{Id}_A \quad \text{i.e., } A \in \mathbf{ALG}^*(\text{CL}).$$

On the other hand, show that:

$$\Omega_{\mathbf{4}}(\{a, 1\}) = \text{Id}_A \cup \{\langle 1, a \rangle, \langle 0, \neg a \rangle\} \quad \text{i.e. } \langle \mathbf{4}, \{a, 1\} \rangle \notin \mathbf{MOD}^*(\text{CL}).$$

BCI: recall the algebra M defined via:

\rightarrow^M	\top	t	f	\perp
\top	\top	\perp	\perp	\perp
t	\top	t	f	\perp
f	\top	\perp	t	\perp
\perp	\top	\top	\top	\top

Show that:

$$\Omega_M(\{t, \top\}) = \Omega_M(\{t, f, \top\}) = \text{Id}_M \quad \text{i.e. } M \in \mathbf{ALG}^*(\text{BCI}).$$

Factorizing matrices – 1

Let us take $\mathbf{A} = \langle \mathbf{A}, F \rangle \in \mathbf{MOD}(\mathbf{L})$. We write:

- \mathbf{A}^* for $\mathbf{A}/\Omega_{\mathbf{A}}(F)$
- $[\cdot]_F$ for the canonical epimorphism of \mathbf{A} onto \mathbf{A}^* defined as:

$$[a]_F = \{b \in A \mid \langle a, b \rangle \in \Omega_{\mathbf{A}}(F)\}$$

- \mathbf{A}^* for $\langle \mathbf{A}^*, [F]_F \rangle$.

Lemma 2.4

Let $\mathbf{A} = \langle \mathbf{A}, F \rangle \in \mathbf{MOD}(\mathbf{L})$ and $a, b \in A$. Then:

- 1 $a \in F$ iff $[a]_F \in [F]_F$.
- 2 $\mathbf{A}^* \in \mathbf{MOD}(\mathbf{L})$.
- 3 $[a]_F \leq_{\mathbf{A}^*} [b]_F$ iff $a \rightarrow^{\mathbf{A}} b \in F$.
- 4 $\mathbf{A}^* \in \mathbf{MOD}^*(\mathbf{L})$.

Proof.

- 1 One direction is trivial. Conversely: $[a]_F \in [F]_F$ implies that $[a]_F = [b]_F$ for some $b \in F$; thus $\langle a, b \rangle \in \Omega_A(F)$ and, since $\Omega_A(F)$ is a logical congruence, we obtain $a \in F$.
- 2 Recall that the second claim of Lemma 1.12 says that for a surjective $g: \mathbf{A} \rightarrow \mathbf{B}$ and $F \in \mathcal{F}i_L(\mathbf{A})$ we get $g[F] \in \mathcal{F}i_L(\mathbf{B})$, whenever $g(x) \in g[F]$ implies $x \in F$.
- 3 $[a]_F \leq_{\mathbf{A}^*} [b]_F$ iff $[a]_F \rightarrow^{\mathbf{A}^*} [b]_F \in [F]_F$ iff $[a \rightarrow^{\mathbf{A}} b]_F \in [F]_F$ iff $a \rightarrow^{\mathbf{A}} b \in F$.
- 4 Assume that $\langle [a]_F, [b]_F \rangle \in \Omega_{\mathbf{A}^*}([F]_F)$, i.e., $[a]_F \leq_{\mathbf{A}^*} [b]_F$ and $[b]_F \leq_{\mathbf{A}^*} [a]_F$. Therefore $a \rightarrow^{\mathbf{A}} b \in F$ and $b \rightarrow^{\mathbf{A}} a \in F$, i.e., $\langle a, b \rangle \in \Omega_A(F)$. Thus $[a]_F = [b]_F$. □

Lindenbaum–Tarski matrix

Let L be a weakly implicative logic in \mathcal{L} and $T \in Th(L)$. For every formula φ , we define the set

$$[\varphi]_T = \{\psi \in Fm_{\mathcal{L}} \mid \varphi \leftrightarrow \psi \subseteq T\}.$$

The **Lindenbaum–Tarski matrix** with respect to L and T , \mathbf{LindT}_T , has the filter $\{[\varphi]_T \mid \varphi \in T\}$ and algebraic reduct with the domain $\{[\varphi]_T \mid \varphi \in Fm_{\mathcal{L}}\}$ and operations:

$$c^{\mathbf{LindT}_T}([\varphi_1]_T, \dots, [\varphi_n]_T) = [c(\varphi_1, \dots, \varphi_n)]_T$$

Clearly, for every $T \in Th(L)$ we have:

$$\mathbf{LindT}_T = \langle Fm_{\mathcal{L}}, T \rangle^*.$$

The second completeness theorem

Theorem 2.5

Let L be a weakly implicative logic. Then for any set Γ of formulae and any formula φ the following holds:

$$\Gamma \vdash_L \varphi \quad \text{iff} \quad \Gamma \models_{\text{MOD}^*(L)} \varphi.$$

Proof.

Using just the soundness part of the first completeness theorem it remains to prove:

$$\Gamma \models_{\text{MOD}^*(L)} \varphi \quad \text{implies} \quad \Gamma \vdash_L \varphi.$$

Take Lindenbaum–Tarski matrix $\mathbf{LindT}_{\text{Th}_L(\Gamma)} = \langle \mathbf{Fm}_{\mathcal{L}}, \text{Th}_L(\Gamma) \rangle^*$ and evaluation $e(\psi) = [\psi]_{\text{Th}_L(\Gamma)}$. As clearly $e[\Gamma] \subseteq e[\text{Th}_L(\Gamma)] = [\text{Th}_L(\Gamma)]_{\text{Th}_L(\Gamma)}$, then, as $\mathbf{LindT}_{\text{Th}_L(\Gamma)}$ is an L -model, we have: $e(\varphi) = [\varphi]_{\text{Th}_L(\Gamma)} \in [\text{Th}_L(\Gamma)]_{\text{Th}_L(\Gamma)}$, and so $\varphi \in \text{Th}_L(\Gamma)$ i.e., $\Gamma \vdash_L \varphi$. □

Closure systems and closure operators – 1

Closure system over a set A : a collection of subsets $\mathcal{C} \subseteq \mathcal{P}(A)$ closed under arbitrary intersections and such that $A \in \mathcal{C}$. The elements of \mathcal{C} are called **closed sets**.

Closure operator over a set A : a mapping $C: \mathcal{P}(A) \rightarrow \mathcal{P}(A)$ such that for every $X, Y \subseteq A$:

- 1 $X \subseteq C(X)$,
- 2 $C(X) = C(C(X))$, and
- 3 if $X \subseteq Y$, then $C(X) \subseteq C(Y)$.

Exercise 4

If C is a closure operator, $\{X \subseteq A \mid C(X) = X\}$ is a closure system.

If \mathcal{C} is closure system, $C(X) = \bigcap \{Y \in \mathcal{C} \mid X \subseteq Y\}$ is a closure operator.

Closure systems and closure operators – 2

A closure operator C is **finitary** if for every $X \subseteq A$,
$$C(X) = \bigcup \{C(Y) \mid Y \subseteq X \text{ and } Y \text{ is finite}\}.$$

A closure system \mathcal{C} is called **inductive** if it is closed under unions of upwards directed families (i.e. families $\mathcal{D} \neq \emptyset$ such that for every $A, B \in \mathcal{D}$, there is $C \in \mathcal{D}$ such that $A \cup B \subseteq C$).

Theorem 2.6 (Schmidt Theorem)

A closure operator C is finitary if, and only if, its associated closure system \mathcal{C} is inductive.

Closure systems and closure operators – 3

Each logic L determines a closure system $\text{Th}(L)$ and a closure operator Th_L .

Conversely, given a **structural** closure operator C over $Fm_{\mathcal{L}}$ (for every σ , if $\varphi \in C(\Gamma)$, then $\sigma(\varphi) \in C(\sigma[\Gamma])$), there is a logic L such that $C = \text{Th}_L$.

L is a finitary logic iff Th_L is a finitary closure operator.

The set of all L -filters over a given algebra A , $\mathcal{F}i_L(A)$ is a closure system over A . Its associated closure operator is Fi_L^A .

Corollary 2.7

Given a logic L in a language \mathcal{L} , the following conditions are equivalent:

- 1 L is finitary.
- 2 Fi_L^A is a finitary closure operator for any \mathcal{L} -algebra A .
- 3 $\mathcal{F}i_L(A)$ is an inductive closure system for any \mathcal{L} -algebra A .

Closure systems and closure operators – 4

A **base** of a closure system \mathcal{C} over A is any $\mathcal{B} \subseteq \mathcal{C}$ satisfying one of the following equivalent conditions:

- 1 \mathcal{C} is the coarsest closure system containing \mathcal{B} .
- 2 For every $T \in \mathcal{C} \setminus \{A\}$, there is a $\mathcal{D} \subseteq \mathcal{B}$ such that $T = \bigcap \mathcal{D}$.
- 3 For every $T \in \mathcal{C} \setminus \{A\}$, $T = \bigcap \{B \in \mathcal{B} \mid T \subseteq B\}$.
- 4 For every $Y \in \mathcal{C}$ and $a \in A \setminus Y$ there is $Z \in \mathcal{B}$ such that $Y \subseteq Z$ and $a \notin Z$.

Exercise 5

Show that the four definitions are equivalent.

An element X of a closure system \mathcal{C} over A is called (**finitely**) **\cap -irreducible** if for each (finite non-empty) set $\mathcal{Y} \subseteq \mathcal{C}$ such that $X = \bigcap_{Y \in \mathcal{Y}} Y$, there is $Y \in \mathcal{Y}$ such that $X = Y$.

Abstract Lindenbaum Lemma

An element X of a closure system \mathcal{C} over A is called **maximal w.r.t. an element a** if it is a maximal element of the set $\{Y \in \mathcal{C} \mid a \notin Y\}$ w.r.t. the order given by inclusion.

Proposition 2.8

Let \mathcal{C} be a closure system over a set A and $T \in \mathcal{C}$. Then, T is maximal w.r.t. an element if, and only if, T is \cap -irreducible.

Lemma 2.9

*Let C be a finitary closure operator and \mathcal{C} its corresponding closure system. If $T \in \mathcal{C}$ and $a \notin T$, then there is $T' \in \mathcal{C}$ such that $T \subseteq T'$ and T' is maximal with respect to a . **\cap -irreducible closed sets form a base.***

Operations on matrices – 1

$\langle A, F \rangle$: first-order structure in the equality-free predicate language with function symbols from \mathcal{L} and a unique unary predicate symbol interpreted by F .

Submatrix: $\langle A, F \rangle \subseteq \langle B, G \rangle$ if $A \subseteq B$ and $F = A \cap G$. Operator: $\mathbf{S}(\langle A, F \rangle)$ is the class of all subalgebras of $\langle A, F \rangle$.

Homomorphic image: $\langle B, G \rangle$ is a homomorphic image of $\langle A, F \rangle$ if it exists $h: A \rightarrow B$ homomorphism of algebras such that $h[F] \subseteq G$. Operator \mathbf{H} .

Strict homomorphic image: $\langle B, G \rangle$ is a strict homomorphic image of $\langle A, F \rangle$ if it exists $h: A \rightarrow B$ homomorphism of algebras such that $h[F] \subseteq G$ and $h[A \setminus F] \subseteq B \setminus G$. Operator \mathbf{H}_S .

Isomorphic image: Image by a bijective strict homomorphism. Operator \mathbf{I} .

Direct product: Given matrices $\{\langle A_i, F_i \rangle \mid i \in I\}$, their direct product is $\langle A, F \rangle$, where $A = \prod_{i \in I} A_i$,
 $f^A(a_1, \dots, a_n)(i) = f^{A_i}(a_1(i), \dots, a_n(i))$. $F = \prod_{i \in I} F_i$. $\pi_j : A \rightarrow A_j$.
Operator **P**.

Exercise 6

Let L be a weakly implicative logic. Then:

- 1 $\mathbf{SP(MOD(L))} \subseteq \mathbf{MOD(L)}$.
- 2 $\mathbf{SP(MOD^*(L))} \subseteq \mathbf{MOD^*(L)}$.

Subdirect products and subdirect irreducibility

A matrix \mathbf{A} is said to be **representable as a subdirect product** of the family of matrices $\{\mathbf{A}_i \mid i \in I\}$ if there is an embedding homomorphism α from \mathbf{A} into the direct product $\prod_{i \in I} \mathbf{A}_i$ such that for every $i \in I$, the composition of α with the i -th projection, $\pi_i \circ \alpha$, is a surjective homomorphism. In this case, α is called a **subdirect representation**, and it is called **finite** if I is finite.

Operator $\mathbf{P}_{\text{SD}}(\mathbb{K})$.

A matrix $\mathbf{A} \in \mathbb{K}$ is **(finitely) subdirectly irreducible relative to \mathbb{K}** if for every (finite non-empty) subdirect representation α of \mathbf{A} with a family $\{\mathbf{A}_i \mid i \in I\} \subseteq \mathbb{K}$ there is $i \in I$ such that $\pi_i \circ \alpha$ is an isomorphism. The class of all (finitely) subdirectly irreducible matrices relative to \mathbb{K} is denoted as $\mathbb{K}_{\text{R(F)SI}}$.

$$\mathbb{K}_{\text{RSI}} \subseteq \mathbb{K}_{\text{RFSI}}.$$

Theorem 2.10

Given a weakly implicative logic L and $\mathbf{A} = \langle A, F \rangle \in \mathbf{MOD}^*(L)$, we have:

- 1 $\mathbf{A} \in \mathbf{MOD}^*(L)_{\text{RSI}}$ iff F is \cap -irreducible in $\mathcal{Fi}_L(A)$.
- 2 $\mathbf{A} \in \mathbf{MOD}^*(L)_{\text{RFSI}}$ iff F is finitely \cap -irreducible in $\mathcal{Fi}_L(A)$.

Theorem 2.11

If L is a finitary weakly implicative logic, then

$$\mathbf{MOD}^*(L) = \mathbf{P}_{\text{SD}}(\mathbf{MOD}^*(L)_{\text{RSI}}),$$

in particular every matrix in $\mathbf{MOD}^(L)$ is representable as a subdirect product of matrices in $\mathbf{MOD}^*(L)_{\text{RSI}}$.*

The third completeness theorem

Theorem 2.12

Let L be a finitary weakly implicative logic. Then

$$\vdash_L = \models_{\mathbf{MOD}^*(L)_{RSI}}.$$

Leibniz operator: the function giving for each $F \in \mathcal{Fi}_L(\mathbf{A})$ the Leibniz congruence $\Omega_A(F)$.

Proposition 2.13

Let L be a weakly implicative logic L and A an \mathcal{L} -algebra. Then

- 1 Ω_A is monotone: if $F \subseteq G$ then $\Omega_A(F) \subseteq \Omega_A(G)$.
- 2 Ω_A commutes with inverse images by homomorphisms: for every \mathcal{L} -algebra B , homomorphism $h: A \rightarrow B$, and $F \in \mathcal{Fi}_L(B)$:

$$\Omega_A(h^{-1}[F]) = h^{-1}[\Omega_B(F)] = \{\langle a, b \rangle \mid \langle h(a), h(b) \rangle \in \Omega_B(F)\}.$$

- 3 $\Omega_A[\mathcal{Fi}_L(A)] = \mathbf{Con}_{\mathbf{ALG}^*(L)}(A)$.

$\mathbf{Con}_{\mathbf{ALG}^*(L)}(A)$ is the set ordered by inclusion of congruences of A giving a quotient in $\mathbf{ALG}^*(L)$.

Example

Recall that for the algebra $\mathbf{M} \in \mathbf{ALG}^*(\mathbf{BCI})$ defined via:

$\rightarrow^{\mathbf{M}}$	\top	t	f	\perp
\top	\top	\perp	\perp	\perp
t	\top	t	f	\perp
f	\top	\perp	t	\perp
\perp	\top	\top	\top	\top

we have

$$\Omega_{\mathbf{M}}(\{t, \top\}) = \Omega_{\mathbf{M}}(\{t, f, \top\}) = \text{Id}_{\mathbf{M}} \quad \text{i.e., } \Omega_{\mathbf{M}} \text{ is not injective}$$

Theorem 2.14

Given any weakly implicative logic L , TFAE:

- 1 For every \mathcal{L} -algebra A , the Leibniz operator Ω_A is a **lattice isomorphism** from $\mathcal{F}i_L(A)$ to $Con_{\mathbf{ALG}^*(L)}(A)$.
- 2 For every $\langle A, F \rangle \in \mathbf{MOD}^*(L)$, F is the least L -filter on A .
- 3 The Leibniz operator $\Omega_{Fm_{\mathcal{L}}}$ is a **lattice isomorphism** from $\mathbf{Th}(L)$ to $Con_{\mathbf{ALG}^*(L)}(Fm_{\mathcal{L}})$.
- 4 There is a set of equations \mathcal{T} in one variable such that
(Alg) $p \dashv\vdash_L \{ \mu(p) \leftrightarrow \nu(p) \mid \mu \approx \nu \in \mathcal{T} \}$.
- 5 There is a set of equations \mathcal{T} in one variable such that for each $\mathbf{A} = \langle A, F \rangle \in \mathbf{MOD}^*(L)$ and each $a \in A$ holds: $a \in F$ if, and only if, $\mu^A(a) = \nu^A(a)$ for every $\mu \approx \nu \in \mathcal{T}$.

In the last two items the sets \mathcal{T} can be taken the same.

Definition 2.15

We say that a logic L is **algebraically implicative** if it is weakly implicative and satisfies one of the equivalent conditions from the previous theorem.

In this case, $\mathbf{ALG}^*(L)$ is called an **equivalent algebraic semantics** for L and the set \mathcal{T} is called a **truth definition**.

Example 2.16

In many cases, one equation is enough for the truth definition. For instance, in classical logic, intuitionism, t-norm based fuzzy logics, etc. the truth definition is $\{p \approx \bar{1}\}$. Linear logic is algebraically implicative with $\mathcal{T} = \{p \wedge \bar{1} \approx \bar{1}\}$.

Different logics with the same algebras

Exercise 7

$\mathcal{L} = \{\neg, \rightarrow\}$. Algebra A with domain $\{0, \frac{1}{2}, 1\}$ and operations:

	\neg
0	1
$\frac{1}{2}$	$\frac{1}{2}$
1	0

	\rightarrow	0	$\frac{1}{2}$	1
0		1	1	1
$\frac{1}{2}$		$\frac{1}{2}$	1	1
1		0	$\frac{1}{2}$	1

$$\mathbb{L}_3 = \models_{\langle A, \{1\} \rangle}$$

[three-valued Łukasiewicz logic]

$$\mathbb{J}_3 = \models_{\langle A, \{\frac{1}{2}, 1\} \rangle}$$

[Da Costa, D'Ottaviano]

Defined connectives: $\bar{1} = p \rightarrow p$, $\diamond p = \neg p \rightarrow p$

\mathbb{L}_3 and \mathbb{J}_3 are both algebraically implicative with

L	ALG*(L)	$\mathcal{T}(p)$
\mathbb{L}_3	$\mathbf{Q}(A)$	$\{p \approx \bar{1}\}$
\mathbb{J}_3	$\mathbf{Q}(A)$	$\{\diamond p \approx \bar{1}\}$

Equational consequence

An **equation** in the language \mathcal{L} is a formal expression of the form $\varphi \approx \psi$, where $\varphi, \psi \in Fm_{\mathcal{L}}$.

We say that an equation $\varphi \approx \psi$ is a **consequence** of a set of equations Π w.r.t. a class \mathbb{K} of \mathcal{L} -algebras if for each $A \in \mathbb{K}$ and each A -evaluation e we have $e(\varphi) = e(\psi)$ whenever $e(\alpha) = e(\beta)$ for each $\alpha \approx \beta \in \Pi$; we denote it by $\Pi \models_{\mathbb{K}} \varphi \approx \psi$.

Proposition 2.17

Let L be a weakly implicative logic and $\Pi \cup \{\varphi \approx \psi\}$ a set of equations. Then

$$\Pi \models_{\mathbf{ALG}^*(L)} \varphi \approx \psi \quad \text{iff} \quad \{\alpha \leftrightarrow \beta \mid \alpha \approx \beta \in \Pi\} \vdash_L \varphi \leftrightarrow \psi.$$

Alternatively, using translation $\rho[\Pi] = \bigcup_{\alpha \approx \beta \in \Pi} (\alpha \leftrightarrow \beta)$:

$$\Pi \models_{\mathbf{ALG}^*(L)} \varphi \approx \psi \quad \text{iff} \quad \rho[\Pi] \vdash_L \rho(\varphi \approx \psi).$$

Characterizations of algebraically implicative logics

We have defined a translation ρ from (sets of) equations to sets of formulae using \leftrightarrow .

Analogously we define a translation τ from (sets of) formulae to sets of equations using the truth definition \mathcal{T} :

$$\tau[\Gamma] = \{\alpha(\varphi) \approx \beta(\varphi) \mid \varphi \in \Gamma \text{ and } \alpha \approx \beta \in \mathcal{T}\}$$

Theorem 2.18

Given any weakly implicative logic L , TFAE:

- 1 L is algebraically implicative with the truth definition \mathcal{T} .
- 2 There is a set of equations \mathcal{T} in one variable such that:
 - 1 $\Pi \models_{\text{ALG}^*(L)} \varphi \approx \psi$ iff $\rho[\Pi] \vdash_L \rho(\varphi \approx \psi)$
 - 2 $p \not\vdash_L \rho[\tau(p)]$
- 3 There is a set of equations \mathcal{T} in one variable such that:
 - 1 $\Gamma \vdash_L \varphi$ iff $\tau[\Gamma] \models_{\text{ALG}^*(L)} \tau(\varphi)$
 - 2 $p \approx q \models_{\text{ALG}^*(L)} \tau[\rho(p \approx q)]$

Finitary algebraically implicative logics and quasivarieties

A quasivariety is a class of algebras described by quasiequations, formal expressions of the form

$\bigwedge_{i=1}^n \alpha_i \approx \beta_i \Rightarrow \varphi \approx \psi$, where $\alpha_1, \dots, \alpha_n, \beta_1, \dots, \beta_n, \varphi, \psi \in Fm_{\mathcal{L}}$.

Proposition 2.19

If \mathbf{L} is a finitary algebraically implicative logic, then it has a finite truth definition and $\mathbf{ALG}^(\mathbf{L})$ is a quasivariety.*

Rasiowa-implicative and regularly implicative logics

Definition 2.20

We say that a weakly implicative logic L is

- **regularly implicative** if:

$$\text{(Reg)} \quad \varphi, \psi \vdash_L \psi \rightarrow \varphi.$$

- **Rasiowa-implicative** if:

$$\text{(W)} \quad \varphi \vdash_L \psi \rightarrow \varphi.$$

Proposition 2.21

A weakly implicative logic L is regularly implicative iff all the filters of the matrices in $\mathbf{MOD}^(L)$ are singletons.*

Proposition 2.22

A regularly implicative logic L is Rasiowa-implicative iff for each $\mathbf{A} = \langle \mathbf{A}, \{t\} \rangle \in \mathbf{MOD}^(L)$ the element t is the maximum of $\leq_{\mathbf{A}}$.*

Proposition 2.23

Each Rasiowa-implicative logic is regularly implicative and each regularly implicative logic is algebraically implicative.

The following logics are Rasiowa-implicative:

- classical logic
- global modal logics
- intuitionistic and superintuitionistic logics
- many fuzzy logics (Łukasiewicz, Gödel-Dummett, product logics, HL, MTL, ...)
- substructural logics with weakening
- inconsistent logic
- ...

Example 2.24

- The equivalence fragment of classical logic is a regularly implicative but not Rasiowa-implicative logic.
- Linear logic is algebraically, but not regularly, implicative.
- The logic BCI is weakly, but not algebraically, implicative.