

Nonlinearity in Normal Human EEG: Cycles, temporal asymmetry, nonstationarity and randomness, not chaos

Milan Paluš

Santa Fe Institute, 1399 Hyde Park Road, Santa Fe, NM 87501, USA

and

Institute of Computer Science, Academy of Sciences of the Czech Republic

Pod vodárenskou věží 2, 182 07 PRAGUE 8, Czech Republic

E-mail: mp@uivt.cas.cz, mp@santafe.edu

November 7, 1995

Abstract

Two-hour vigilance and sleep EEG recordings from five healthy volunteers were analyzed using a method for identifying nonlinearity and chaos, which combines the redundancy – linear redundancy approach with the surrogate data technique. A nonlinear component in the EEG was detected, however, inconsistent with the hypothesis of low-dimensional chaos. A possibility, that a temporally asymmetric process may underlie or influence the EEG dynamics, was indicated. A process, that merges nonstationary nonlinear deterministic oscillations with randomness, is proposed for an explanation of observed properties of the analyzed EEG signals. Taking these results into consideration, the use of dimensional and related chaos-based algorithms in quantitative EEG analysis is critically discussed.

1 Introduction

During the last decade there has been a sustained interest in describing neural processes and brain-signals, especially the electroencephalogram (EEG), within the context of nonlinear dynamics and theory of deterministic chaos (see Rapp et al. 1989, Başar 1990, Jansen 1991, Freeman 1992, for reviews). If the nature of analyzed signals was actually low-dimensional, the published results could be of immense importance for theoretical neuroscience and neurological and psychiatric clinical practice. However, confidence of results obtained from experimental data, such as finite dimensions or positive Lyapunov exponents, and reliability of chaos-based algorithms in general, have recently come under question, and alternative methods for identifying possible nonlinear determinism in experimental time series have been proposed (see Weigend & Gershenfeld 1993, Paluš 1995, and references therein).

This paper presents results from applying a method for identifying nonlinearity and chaos, which combines the redundancy – linear redundancy approach with the surrogate data technique, to vigilance and sleep EEG recordings of approximately two-hour duration from five healthy volunteers. The method consists of a test for nonlinearity, in which a null hypothesis of a linear stochastic process is tested by statistics based on nonlinear and linear redundancies (Paluš 1995). The presence of a signature of chaos, positive Kolmogorov-Sinai entropy (KSE), is assessed using the marginal redundancy approach (Paluš 1993). Employing these methods (briefly introduced in Sec. 2, for details we refer to Paluš 1993, 1995, 1994a), a nonlinear component was detected in all analyzed EEG records, however, signatures of low-dimensional chaos were absent. Similar results have been published by Theiler & Rapp (1995), Prichard & Theiler (1994), Theiler et al. (1992) and Casdagli (1992). These authors rejected low-dimensional chaos, however, in some cases they also reported very weak evidence for nonlinearity. In this study very long EEG recordings were analyzed, which could be

the source of greater statistical power of the performed tests and thus the stronger evidence for nonlinearity, than in the other studies. We also indicate that the EEG may possess temporal asymmetry as a nonlinear property.

A process, that merges nonstationary nonlinear deterministic oscillations with randomness, is proposed for an explanation of observed properties of the analyzed EEG signals. Taking these results into consideration, the use of dimensional and related chaos-based algorithms in quantitative EEG analysis is critically discussed.

2 Principles of the Methods

Consider n discrete random variables X_1, \dots, X_n with sets of values Ξ_1, \dots, Ξ_n , respectively. The probability distribution for an individual X_i is $p(x_i) = \Pr\{X_i = x_i\}$, $x_i \in \Xi_i$. We denote the probability distribution function by $p(x_i)$, rather than $p_{X_i}(x_i)$, for convenience. Analogously, the joint distribution for the n variables X_1, \dots, X_n is $p(x_1, \dots, x_n)$. The redundancy $R(X_1; \dots; X_n)$, in the case of two variables also known as mutual information $I(X_1; X_2)$, quantifies average amount of common information, contained in the n variables X_1, \dots, X_n :

$$R(X_1; \dots; X_n) = \sum_{x_1 \in \Xi_1} \dots \sum_{x_n \in \Xi_n} p(x_1, \dots, x_n) \log \frac{p(x_1, \dots, x_n)}{p(x_1) \dots p(x_n)}. \quad (1)$$

The marginal redundancy $\varrho(X_1, \dots, X_{n-1}; X_n)$ quantifies the average amount of information about the variable X_n , contained in the $n - 1$ variables X_1, \dots, X_{n-1} :

$$\varrho(X_1, \dots, X_{n-1}; X_n) = \sum_{x_1 \in \Xi_1} \dots \sum_{x_n \in \Xi_n} p(x_1, \dots, x_n) \log \frac{p(x_1, \dots, x_n)}{p(x_1, \dots, x_{n-1})p(x_n)}. \quad (2)$$

The relation

$$\varrho(X_1, \dots, X_{n-1}; X_n) = R(X_1; \dots; X_n) - R(X_1; \dots; X_{n-1}) \quad (3)$$

can be derived by simple manipulation.

When the discrete variables X_1, \dots, X_n are obtained from continuous variables on a continuous probability space, then the redundancies depend on a partition ξ chosen to discretize the space. Various strategies have been proposed to define an optimal partition for estimating redundancies of continuous variables (see Paluš 1995, Weigend & Gershenfeld 1993, and references therein). Here we use the “marginal equiquantization” method described in detail in (Paluš 1993, 1995). Pompe (1993) also describes this kind of partitioning.

Now, let the n variables X_1, \dots, X_n have zero means, unit variances and correlation matrix \mathbf{C} . Then, we define the *linear redundancy* $L(X_1; \dots; X_n)$ of X_1, X_2, \dots, X_n as

$$L(X_1; \dots; X_n) = -\frac{1}{2} \sum_{i=1}^n \log(\sigma_i), \quad (4)$$

where σ_i are the eigenvalues of the $n \times n$ correlation matrix \mathbf{C} .

If X_1, \dots, X_n have an n -dimensional Gaussian distribution, then $L(X_1; \dots; X_n)$ and $R(X_1; \dots; X_n)$ are theoretically equivalent (Morgera 1985).

Based on (3) we define the *linear marginal redundancy* $\lambda(X_1, \dots, X_{n-1}; X_n)$, quantifying linear dependence of X_n on X_1, \dots, X_{n-1} , as

$$\lambda(X_1, \dots, X_{n-1}; X_n) = L(X_1; \dots; X_n) - L(X_1; \dots; X_{n-1}). \quad (5)$$

In practical applications one deals with a time series $\{y(t)\}$, considered as a realization of a stochastic process $\{Y(t)\}$, which is stationary and ergodic. Then, due to ergodicity, all the subsequent information-theoretic functionals are estimated using time averages instead of ensemble averages, and the variables X_i are substituted as

$$X_i = y(t + (i - 1)\tau). \quad (6)$$

Due to stationarity the redundancies

$$R^n(\tau) \equiv R(y(t); y(t + \tau); \dots; y(t + (n - 1)\tau)), \quad (7)$$

$$\varrho^n(\tau) \equiv \varrho(y(t), y(t + \tau), \dots, y(t + (n - 2)\tau); y(t + (n - 1)\tau)), \quad (8)$$

$$L^n(\tau) \equiv L(y(t); y(t + \tau); \dots; y(t + (n - 1)\tau)), \quad (9)$$

$$\lambda^n(\tau) \equiv \lambda(y(t), y(t + \tau), \dots, y(t + (n - 2)\tau); y(t + (n - 1)\tau)) \quad (10)$$

are functions of n and τ , independent of t .

The surrogate-data based nonlinearity tests consist of computing a *nonlinear* statistic from data under study and from an ensemble of realizations of a linear stochastic process, which mimics “linear properties” of the studied data. If the computed statistic for the original data is significantly different from the values obtained for the surrogate set, one can infer that the data were not generated by a linear process; otherwise the null hypothesis, that a linear model fully explains the data, is accepted. For the purpose of such test the surrogate data must preserve the spectrum¹ and consequently, the autocorrelation function of the series under study. An isospectral linear stochastic process to a series can be constructed by computing the Fourier transform (FT) of the series, keeping unchanged the magnitudes of the Fourier coefficients, but randomizing their phases and computing the inverse FT into the time domain. Different realizations of the process are obtained using different sets of the random phases.

To evaluate the test, the test statistic is defined as the difference between the redundancy obtained for the original data and the mean redundancy of a set of surrogates, in the number of standard deviations (SD’s) of the latter. Thus both the redundancies and redundancy-based statistics can be evaluated as functions of the lag τ and the embedding dimension n .

The redundancy $R^n(\tau)$ (or, $\varrho^n(\tau)$), based on probability distributions, measures general dependences among the series $\{y(t)\}$ and its lagged versions, whereas the linear redundancy $L^n(\tau)$ (or, $\lambda^n(\tau)$), based on correlations, reflects only their linear relations. Comparing the plots of $R^n(\tau)$ and $L^n(\tau)$ (or $\varrho^n(\tau)$ and $\lambda^n(\tau)$) can provide an informal test for important nonlinearities in the studied data (Paluš et al. 1993). This approach, in (Paluš 1995) referred to as *qualitative testing*, or, *qualitative comparison*, can bring some additional information to the results of the quantitative (surrogate data based) test. Moreover, one can not always construct good surrogate data. That is, despite theoretical expectations, in numerical practice linear properties of the surrogates may differ from those of the data under study. Changes in linear properties are reflected in nonlinear statistics as well, and thus may result in spurious detection of nonlinearity in linear data (Paluš 1995). Therefore, we also evaluate the statistic based on the linear redundancy $L^n(\tau)$, which specifically reflects changes in linear properties. Then, only those significant differences in the nonlinear statistic can reliably count for nonlinearity, which are not detected in the linear statistic (Paluš 1995).

If a time series $\{y(t)\}$ was generated by an m -dimensional chaotic system, then for $n > m$ and under some conditions the marginal redundancy $\varrho^n(\tau)$ decreases with τ as

$$\varrho^n(\tau) \approx H_1 - |\tau|h, \quad (11)$$

where h is the Kolmogorov-Sinai entropy (KSE) of the underlying dynamical system, and H_1 is a constant related to precision (quantization) of data (Fraser 1989, Paluš 1993). Fulfilling the conditions for (11) can be limited by an available amount of data (Paluš 1994a, Paluš submitted). Therefore we propose a comparison of the slopes of the redundancy $\varrho^n(\tau)$ and the linear redundancy $\lambda^n(\tau)$ (or, the slopes of $\varrho^n(\tau)$ obtained from the data under study and $\varrho^n(\tau)$ from its surrogates) to find whether the decrease of $\varrho^n(\tau)$ is a nonlinear

¹Also, preservation of histogram is usually required. A histogram transformation used for this purpose is described in (Paluš 1995) and references within.

(possibly chaotic) phenomenon, or can be explained by a linear stochastic process. The latter case renders a hypothesis of deterministic chaos improbable.

3 Material and Methods

Five paid, physically and psychically healthy volunteers (see Tab. 1), chosen from a larger group by passing standard EEG-study criteria, participated in the experiment after signing the informed consent. The five subjects were recorded in two different states:

1. Relaxed vigilance with closed eyes. Vigilance was assured by the volunteers' participation in a simple mental task: Subjects were asked to listen to a text recorded on a tape-recorder and to count occurrences of the conjunction "but". (For brevity, the term "vigilant state" will be used henceforth to refer to this state.)
2. Sleep evoked by a 1 or 2 mg dose (according to individual sensitivity) of flunitrazepam (tablets). (Henceforth referred to as "sleep state".)

Eight channels of EEG data, sampled by 256-Hz frequency and 12-bit precision, were recorded from the positions O_1 , O_2 , T_5 , T_6 , F_7 , F_8 , P_z and F_z , using the common Goldman reference electrode. Each record consists of ninety 90-second segments ($90 \times 23,040$ samples).

The marginal redundancies $\lambda^n(\tau)$ and $\varrho^n(\tau)$ and the related linear and nonlinear statistics were calculated from the redundancies $L^n(\tau)$ and $R^n(\tau)$, estimated from the data and the surrogates (30 realizations of the surrogates for each dataset). Due to the segmentation of the data, the n-tuples $y(t); y(t + \tau); \dots; y(t + (n - 1)\tau)$ were constructed only inside the individual segments, whereas final probability distributions and the correlation matrices were averaged through all the segments. For the qualitative comparisons, the original-data redundancies $R^n(\tau)[o]$ and $L^n(\tau)[o]$ were computed from entire 23,040-sample segments, which gives the effective series length $N = 2,037,600$, given embedding dimensions $n = 2 - 5$, a lag range 1 - 100 and the segmentation. In the surrogate data tests 16,384 (16k)-sample subsegments were used, employing fast Fourier transform (Press et al. 1986), requiring a series length equal to a power of two. This yields the effective series length $N = 1,438,560$, applicable for the redundancy estimations. The surrogate data were generated for each segment separately, i.e., using 16,384 samples. Gaussianization (Paluš 1995) was performed in the quantitative analysis; before the analysis, the data were filtered using a notch filter in the frequency domain, in order to eliminate interference from the recording equipment.

All the redundancies were computed for embedding dimensions $n = 2 - 5$ using $Q = 4 - 16$ marginal equiquantal (equipopulated) bins. In the qualitative testing, the redundancies were also computed with Q up to 64, but n was only 2 and 3.

4 Results

Figure 1 presents the analysis results of the EEG of Subj. 2 recorded in position O_1 in the vigilant state. In the qualitative comparison there are practically no differences between the linear marginal redundancy $\lambda^n(\tau)$ (Fig. 1a) and the marginal redundancy $\varrho^n(\tau)$ (Fig. 1b). The result of quantitative analysis is different: There are several highly significant differences (about 10 SD's) detected in the statistic based on nonlinear redundancy (Fig. 1d). The possibility that these significances are spurious (caused by imperfect surrogates) is ruled out by the results in the statistic based on linear redundancy, where no significant differences (differences between -1 and 1 SD) were found (Fig. 1c). The hypothesis of a linear stochastic process was thus rejected.

Looking at the nonlinear redundancy statistic (Fig. 1d) in detail, one can see that the significant differences are periodically distributed and located in the lags in which $\varrho^n(\tau)$ has its minima. This means that $\varrho^n(\tau)$ is *significantly* greater than zero also in its minima, which is not apparent from Fig. 1b, esp. in the case of $\varrho^2(\tau)$. Thus, $\varrho^2(\tau) > 0$ and the variables $x(t)$ and $x(t + \tau)$ are dynamically (deterministically) connected for the whole range of considered lags, i.e., up to 100 samples, or, 390 msec. In the case of the surrogates, $\varrho^2(\tau)$ vanishes in its minima. For other lags the differences between the EEG data and the surrogates are




Figure 1: (a) Linear marginal redundancy and (b) marginal redundancy, as functions of time lag, for the vigilance EEG of Subj. 2 recorded from position O_1 . The four different curves in each figure are the redundancies for different embedding dimensions, $n = 2 - 5$, reading from bottom to top. (c) Linear redundancy statistic, and (d) nonlinear redundancy statistic, as functions of time lag and for embedding dimensions $n = 2 - 5$, for the same EEG record.

not significant. This phenomenon, typical for *nonlinear deterministic oscillations* (Paluš 1995), will be called *periodic nonlinearity*.

The long-term decreasing trend in $\varrho^n(\tau)$ is exactly reflected also in linear $\lambda^n(\tau)$; and is fully explained by the isospectral linear stochastic process, because there is no indication of a decreasing pattern in the time-lag dependence of differences (Fig. 1d; cf. Fig. 3d in Paluš 1994a).

The results in Figs. 1b and 1d were obtained using $Q = 4$ marginal equiquantal bins. The tests using finer partitions, possible due to large amount of available data, were also performed (see Sec. 3). Using finer partitions, however, did not bring any new information either in the qualitative comparison, nor in the quantitative analysis (Paluš 1994b). This phenomenon is typical for all EEG data analyzed.

Results similar to those depicted in Fig. 1 and described above, were obtained from the EEG signals of all five subjects, recorded in the vigilant state from positions O_1 , O_2 , T_5 , T_6 , P_z and F_z , i.e., from all channels except the two frontal ones, F_7 and F_8 . For the sleep state and the above six channels we can also present one typical example. The analysis results obtained from the EEG of Subj. 1, recorded in the sleep state from position O_1 , are presented in Fig. 2. In general, the sleep EEG's are less coherent than the vigilance ones, the values of the redundancies are about ten-times smaller and decrease faster, i.e., the coherence length² is shorter. The results of the nonlinearity tests, however, are the same as those for the vigilance EEG, described above: No qualitative differences between $\varrho^n(\tau)$ computed from the EEG and from the surrogates³ were found (Figs. 2a,b). No significant differences (from -1 to 1.6 SD's) were found in the linear redundancy statistic (Fig. 2c); but several highly significant differences (tens of SD's) were detected in the nonlinear redundancy statistic (Fig. 2d). These significant differences are located in the lags in which $\varrho^n(\tau)$ has its minima.

Properties of the EEG signals recorded in the frontal loci (F_7 and F_8 ; note that F_z was included in the above 6 channels) in both the vigilant and sleep states are similar to those obtained from other channels in the sleep state, with two differences:

1. In addition to the periodic nonlinearity, significant differences in "short" lags, i.e., in the lags 1 – 2 samples (4 – 8 ms) or 1 – 3 samples (4 – 12 ms) were detected (Fig. 3b). We will henceforth refer to this phenomenon as "short-lag nonlinearity".
2. In three subjects in both the vigilant and sleep states, this short-lag nonlinearity was the only significant difference detected (Fig. 3a). We observed, however, that the spectra of these records are dominated by slow activity⁴. When this slow activity was eliminated by a digital filter in spectral domain, a typical pattern of periodic nonlinearity emerged. The short-lag nonlinearity, however, was not eliminated.

An example of this phenomenon is presented in Fig. 3. The analysis results of the vigilant-state EEG of Subj. 1, recorded in position F_7 , are depicted in Figs. 3a, c. (We do not present $\varrho^n(\tau)$ from the surrogates, which is not qualitatively different from $\varrho^n(\tau)$ obtained from the EEG, nor the linear redundancy statistic, in which no significant differences were found.) In the nonlinear statistic the only significant differences are located in the lags 1 – 2 samples (Fig. 3a). The lag-dependence of $\varrho^n(\tau)$ (Fig. 3c), however, is similar to that presented in Fig. 2b, showing the typical sleep-state result. After eliminating the slow activity from the signal, the lag dependence of $\varrho^n(\tau)$ is close to the typical vigilance pattern (Fig. 1), but the coherence length is shorter (Fig. 3d). Results from the quantitative test are also typical, i.e., the significant differences in the minima of $\varrho^n(\tau)$ were detected, but with addition of significant differences in the lags 1 and 2 samples (Fig. 3b).

In general, the properties of EEG signals recorded in frontal positions F_7 and F_8 are not different from the properties of other EEG signals described above. The only difference is the short-lag nonlinearity, which is typical for these channels, but can also be found in different locations (Tab. 1).

Defining qualitative differences between $\varrho^n(\tau)[o]$ from the original data and $\varrho^n(\tau)[s]$ from the surrogates

²Under the coherence length we understand here such time lag γ that $\varrho^2(\tau) > 0$ significantly $\forall \tau \leq \gamma$, where $\gamma, \tau \geq 0$.

³The comparison of $\varrho^n(\tau)$ computed from the EEG data with $\varrho^n(\tau)$ from the surrogates is equivalent to the comparison of $\lambda^n(\tau)$ with $\varrho^n(\tau)$ from the EEG, and may be even more reliable (Paluš 1995). Therefore we will use the former in all remaining tests presented here. $\varrho^n(\tau)$ from the surrogates means the mean values of $\varrho^n(\tau)$ obtained from 30 realizations of the surrogates.

⁴Unfortunately we were not able to determine whether it was slow EEG activity or eye artifacts, but the latter are more probable.




Figure 2: Marginal redundancy for (b) the sleep EEG of Subj. 1 from position O_1 and (a) its surrogates, as functions of the time lag. The four different curves in each figure are the redundancies for different embedding dimensions, $n = 2 - 5$, reading from bottom to top. (c) Linear redundancy statistic, and (d) nonlinear redundancy statistic, as functions of the time lag and for embedding dimensions $n = 2 - 5$, for the same EEG record.




Figure 3: (a, b) Nonlinear redundancy statistic and (c, d) marginal redundancy for the vigilance EEG of Subj. 1 from position F_7 , (a, c) before and (b, d) after high-pass filtering, as functions of time lag and for embedding dimensions $n = 2 - 5$.




Figure 4: (a) Linear $\lambda^2(\tau)$ and (b) nonlinear marginal redundancy $\rho^2(\tau)$ for the sleep EEG of Subj. 5 from position P_z (full lines) and its surrogates (mean value – dashed lines, mean \pm SD – dotted lines). (c, d): The same as (b), but (c) for the vigilance EEG of Subj. 1, position O_1 and (d) for the simulated data with gradually changing fast frequency during oscillatory episodes, discussed in Sec. 5.

Subject	1	2	3	4	5
Age	43	40	44	31	20
Sex	F	F	M	F	M
O_1 – vigil	P	P	P	P	P
O_1 – sleep	P	P	P	P S	P
O_2 – vigil	P	P	P	P	P
O_2 – sleep	P	P	P	P S	P
T_5 – vigil	P S	P S	P	P S	P S
T_5 – sleep	P	P S	P S	P S	P S
T_6 – vigil	P S	P S	P	P S	P S
T_6 – sleep	P	P S	P S	P S	P S
P_z – vigil	P	P	P	P S	P S
P_z – sleep	P	P	P	P S	P
F_z – vigil	P	P	P	P S	P S
F_z – sleep	P	P	P S	P S	P
F_7 – vigil	F S	P S	F S	F S	P S
F_7 – sleep	F S	P S	F S	F S	P S
F_8 – vigil	F S	P S	F S	F S	P S
F_8 – sleep	F S	P S	F S	F S	P S

Table 1: *Basic characteristics of the subjects and results of the analysis. Sex: F – female, M – male. Location – state : vigil – the vigilant state, sleep – the sleep state. Analysis results: P – periodic nonlinearity, F – periodic nonlinearity detected after high-pass filtering, S – short-lag nonlinearity.*

is sometimes problematic. Above we have stated that no such differences were found in analysis of the studied EEG data. This is true in the sense of the definition of qualitative differences given in (Paluš 1995), which emphasizes differences in ranges of lags in which $\varrho^n(\tau)$ is positive, locations of extrema and long-term trends of $\varrho^n(\tau)$. Looking at more subtle details, however, one can discover an interesting feature, which differs the nonlinear oscillations in the EEG from trivial periodic nonlinearity, described in (Paluš 1995). For such a detailed comparison, we have plotted in one figure (Fig. 4b) $\varrho^2(\tau)[o]$ from the sleep EEG data of Subj. 5, recorded in position P_z (full line) and $\varrho^2(\tau)[s]$ from its surrogates (mean – dashed line, mean \pm SD – dotted lines). We can observe an asymmetry in the peaks of $\varrho^n(\tau)[o]$ compared with those of $\varrho^n(\tau)[s]$. The fact that this is a nonlinear phenomenon and not an artifact of imperfect surrogates, is demonstrated by comparing $\lambda^2(\tau)[o]$ and $\lambda^2(\tau)[s]$ (Fig. 4a, the same line-type codes as in Fig. 4b): these coincide, thus linear properties of the surrogates are the same as those of the EEG data.

This phenomenon can be found also in other subjects in both the vigilant and sleep states, but in most cases the difference is very small and almost invisible. Another example, when this asymmetry is well detectable, is the vigilant-state EEG from the position O_1 of the Subj. 1 (Fig. 4c). In Fig. 4d this phenomenon is observed in numerically generated data, discussed in Sec. 5.

In summary, all the EEG records possess a nonlinearity. In particular, a type of periodic nonlinearity was detected in all the EEG records (although in some of the EEG’s from frontal channels F_7 and F_8 it was not detected in the raw data, but after high-pass filtering). EEG’s from the frontal channels F_7 and F_8 and from certain other channels also possess the short-lag nonlinearity. These results are summarized in Tab. 1.

5 Discussion of the Results

Periodic nonlinearity, detected in practically all the EEG records analyzed, was characterized by significant differences located in the lags in which the redundancy $\varrho^n(\tau)$ has its minima, while the relations between $y(t)$

and $y(t + \tau)$ in other lags were well reproduced by the linear stochastic surrogates. Also, the linear stochastic surrogates mimic the EEG signals very well from the viewpoint of qualitative comparison. This kind of behavior was observed in periodic processes related to dynamics on limit cycles or tori, while “specifically nonlinear dynamics”, like evolution on strange attractors, was characterized by both qualitative differences and significant results of the quantitative tests in entire ranges of studied lags (Paluš 1994a, 1995). On the other hand, regular periodic processes have zero Kolmogorov-Sinai entropy and thus no long-term decreasing trend in $\varrho^n(\tau)$ (Paluš 1993). Dynamical information loss, measured by a long-term decrease in $\varrho^n(\tau)$, however, is not a phenomenon specific only to chaotic dynamics, but can be caused by any kind of dynamical noise, such as that in autoregressive processes (Paluš 1994a, Paluš submitted), or by a nonstationarity or fluctuation in some parameter of a dynamical system, that might otherwise be regular, i.e., have zero KSE (Paluš 1993). The long-term decreasing trend in $\varrho^n(\tau)$ detected in all EEG records analyzed, has been found consistent with the linear stochastic hypothesis. Thus, studied EEG signals can be a mixture of nonlinear deterministic oscillations, linear stochasticity and/or nonstationarity. In particular, in a numerical study the following simulated time series were constructed:
Episodes of two-frequency oscillations of the form

$$y(t) = A \sin(\omega_1 t) \sin(\omega_2 t), \quad (12)$$

where $\omega_1 = 2\pi/T_1$ represents fast dominant oscillations (e.g., $T_1 = 0.1$ sec.) and $\omega_2 = 2\pi/T_2$ represents a slower “envelope” (e.g., $T_2 = 0.5 - 1.0$ sec.); were randomly distributed in a background of Gaussian noise. Durations of the episodes were random, varying from $0.8T_2$ to $2.2T_2$. Also other parameters were randomized: The amplitude A as $A_t = A_0 + \alpha_t$, the argument of the “envelope sine” as $\omega_2 t + \xi_t$ and the fast frequency ω_1 as $\omega_1(t) = \omega_0 + o_t$, where α_t , ξ_t and o_t are Gaussian deviates with zero means and variances of about 20% of A_0 , $2\pi/3$ and 10% of ω_0 , respectively. Finally, the spectra of these artificial signals are more realistic (i.e., more closely resembling real EEG spectra), when background noise is not white, but filtered into $1/f$ type. Tuning the above parameters (the variances of the random components, the durations of the episodes, the frequency of episode occurrence, and the type of the background noise) one can produce signals, which, when analyzed, yield results practically indistinguishable from the results obtained from various EEG recordings. An example of results from such an artificial time series is presented in Fig. 5: $\varrho^n(\tau)$ of the simulated data (Fig. 5b) and of its surrogates (Fig. 5a), the linear redundancy statistic (Fig. 5c) and the nonlinear redundancy statistic (Fig. 5d). These results are practically the same as those obtained from the vigilance EEG recordings from all but the frontal channels (Fig. 1). By changing the background noise, in particular, by increasing the spectral power of slow frequencies relatively to the power of the main ω_1 -oscillations, the “vigilant-state artificial EEG” can be changed into the “sleep-state artificial EEG”; i.e. the results become similar to those presented in Fig. 2.

When the fast oscillation frequency ω_1 is randomized as described above (or constant) no asymmetries such as those in Fig. 4 are observed. In an additional study we found that such asymmetries may occur when the frequency ω_1 changes gradually during the oscillatory episode. In particular, we used linearly changing period T_1 as $T_1(t) = T_0 + ct$, where t is the time from the beginning of an oscillatory episode. The sign of the parameter c determines on which side of a peak of $\varrho^2(\tau)$ the asymmetry occurs: when c is negative the asymmetry is located on the left side of the peaks, as in the cases presented in Fig. 4; when c is positive, the asymmetry is located on the right side. Thus, if the asymmetries detected in the real EEG data were indeed caused by this mechanism, related c 's are negative and T_1 's decrease, or the dominant frequency (alpha) increases during the oscillatory episodes (alpha spindles).

The above described process is obviously asymmetric with respect to time. Although the actual dynamical mechanisms underlying the EEG can be different⁵, the temporal asymmetry can be detected in the real EEG dynamics⁶. Recently, Chialvo & Millonas (1995) have pointed out the importance of temporally asymmetric fluctuations for biological energy transduction. Possible existence and a role of temporally

⁵A generalized Wiener process with a nonzero drift can be considered as a plausible stochastic alternative to the above deterministic drift in T_1 (ω_1).

⁶The asymmetry in the peaks of $\varrho^2(\tau)$ is not necessarily equivalent to the temporal asymmetry of the underlying process (time series), and the redundancy ϱ^2 (mutual information I) is not a measure suitable for detecting the temporal asymmetry,




Figure 5: Marginal redundancy for (b) the simulated time series discussed in Sec. 5 and (a) its surrogates, as functions of time lag. The four different curves in each figure are the redundancies for different embedding dimensions, $n = 2 - 5$, reading from bottom to top. (c) Linear redundancy statistic, and (d) nonlinear redundancy statistic, as functions of time lag and for embedding dimensions $n = 2 - 5$, for the same simulated series.

asymmetric processes in the brain dynamics could be an interesting subject of research.

The simulated time series, which combine nonstationary deterministic oscillations with randomness, namely, in which oscillatory episodes with some randomized parameters are randomly distributed in a noisy background, can mimic all properties of the real EEG signals analyzed⁷. Of course, this is not evidence that this “model” explains real EEG dynamics, although it seems more probable that the hypothesis of low-dimensional chaos. Without insisting on a particular model or hypothesis, we can conclude, that in the analyzed EEG recordings nonlinearity was detected, however, not consistent with the hypothesis of low-dimensional chaos. Considering this result, applications in computerized EEG analysis of algorithms, designed for characterization of low-dimensional chaotic systems, should be critically reassessed.

Some authors (Mayer-Kress & Layne 1987, Koukkou et al. 1993, Wackermann et al. 1993) no longer insist on interpreting their results, such as finite dimension estimates, as evidence for underlying low-dimensional chaos, but propose these measures, particularly the correlation dimension (CD) (Grassberger & Procaccia 1983), as measures for relative characterization of different EEG recordings. Although these authors demonstrate that such CD estimates can have some discriminating power with respect to EEG’s recorded in different physiological or psychological conditions, considering that the underlying processes can be stochastic, these low numbers, formally obtained from dimensional algorithms, are probably spurious and have no theoretically justified meaning and interpretation. Moreover, it can hardly be established, how robust or sensitive the measures such as CD are, when used in the relative characterization of processes, dimensionality of which can be effectively infinite. For these reasons we do not consider the dimensional and Lyapunov exponents algorithms, based on the particular hypothesis of a low-dimensional attractor, to be promising in the computerized EEG analysis. Nonlinear methods, which can quantify some really existing property of a process under study, should be used. For instance, differences in coherence lengths of various EEG records can induce differences in estimates of CD or Lyapunov exponents. The coherence length of a process is related to its entropy rate and thus can be directly quantified, e.g., by an “approximate entropy” (Pincus et al. 1991), or by “coarse-grained entropy rates” (Paluš submitted), which can be applied to both deterministic-chaotic and stochastic systems, and are designed specifically for noisy, finite-precision and finite-length experimental data.

6 Conclusion

Two-hour vigilance and sleep EEG recordings from five healthy volunteers were analyzed using a method for identifying nonlinearity and chaos, which combines the redundancy – linear redundancy approach with the surrogate data technique. A nonlinear component in the EEG was detected, but was found inconsistent with the hypothesis of low-dimensional chaos. A possibility, that a temporally asymmetric process may underlie or influence the EEG dynamics, was indicated. A process, which merges nonstationary nonlinear deterministic oscillations with randomness, namely, in which deterministic oscillatory episodes are randomly distributed in noisy background, has been proposed for an explanation of observed properties of the analyzed EEG signals. Considering these findings, the use of dimensional and related chaos-based algorithms in the quantitative EEG analysis was criticized, and measures applicable also for nonlinear stochastic processes were proposed instead. However, any EEG study, in which the authors apply nonlinear measures, should also present results of elaborate analysis using linear methods, and/or test hypotheses⁸ using isospectral sur-

due to its symmetry properties ($I(\xi; \eta) = I(\eta; \xi)$). Considering $\xi = y(t)$ and $\eta = y(t + \tau)$, the temporal asymmetry of a series $\{y(t)\}$ can be detected by the significant difference between the probability distributions $p(\xi, \eta)$ and $p(\eta, \xi)$, quantified by the Kullback-Leibler information $KLI = \sum_{\xi} \sum_{\eta} [p(\xi, \eta) - p(\eta, \xi)] \log[p(\xi, \eta)/p(\eta, \xi)]$. Application of this approach led to positive detection of the temporal asymmetry in the EEG, however, yet only a part of the EEG data described in this paper was processed using the KLI measure. Therefore the temporal asymmetry in the EEG is reported here only as a possibility, which requires further study before it is considered a general feature of the EEG.

⁷Except of the short-lag nonlinearity, detected in a part of the datasets (Tab. 1).

⁸Let A and B be two groups of EEG recordings. Positive identification of nonlinearity in individual recordings is not sufficient evidence for the assertion like “The groups A and B can be distinguished only by a nonlinear measure and not by linear methods”. We propose also the comparison of the surrogates of the group A with the surrogates of the group B using the same nonlinear measure as for the comparison of the original data.

rogate data, in order to demonstrate that nonlinear measures are indeed necessary to quantify phenomena under study.

Acknowledgements

The author is grateful to I. Dvořák, I. David, V. Albrecht and G. Mayer-Kress, who made possible to realize this project. D. Chialvo, J. Theiler, D. Prichard, D. Kaplan, B. Hinrichs, J. Klaschka and V. Witkovský are acknowledged for valuable comments and inspiring discussions. Thanks for assistance with computational facilities and data transmission are due to J. Kuriplach, S. Yelich and J. Shannon.

The author was supported by the International Research Fellowship F05 TWO4757 ICP from the National Institutes of Health, the Fogarty International Center, and also by grants to the Santa Fe Institute, including core funding from the John D. and Catherine T. MacArthur Foundation, the National Science Foundation (PHY-8714918), and the U.S. Department of Energy (ER-FG05-88ER25054). An early stage of this study was supported within the framework of the *IBM Academic Initiative in Czecho-Slovakia*, and by the Ministry of Health of the Czech Republic, during the author's stay in the Prague Psychiatric Center, 3rd School of Medicine of the Charles University in Prague.

References

- Başar E. (ed.) (1990) *Chaos in Brain Function*. Springer, Berlin
- Casdagli M (1992) Chaos and deterministic versus stochastic modelling. *JR Stat Soc B* 54: 303-328
- Chialvo DR, Millonas MM (1995) Temporally asymmetric fluctuations are sufficient for the biological energy transduction, Santa Fe Institute Working Paper 95-07-064
- Fraser AM (1989) Information and entropy in strange attractors. *IEEE Transactions on Information Theory* 35: 245-262
- Freeman WJ (1992) Tutorial on Neurobiology: From single neurons to brain chaos. *Int J Bifurcations and Chaos* 2: 451-482
- Grassberger P, Procaccia I (1983) Measuring the strangeness of strange attractors. *Physica D* 9: 189-208
- Jansen BH (1991) Quantitative analysis of electroencephalograms: Is there chaos in the future? *Int J Biomed Comput* 27: 95-123
- Koukkou M, Lehmann D, Wackermann J, Dvořák I, Henggeler B (1993) Dimensional complexity of EEG brain mechanisms in untreated schizophrenia. *Biol Psychiatry* 33: 397-407
- Mayer-Kress G, Layne SP (1987) Dimensionality of the human electroencephalogram. *Annals of the New York Academy of Sciences* 504: 62-86
- Morgera, SD (1985) Information theoretic covariance complexity and its relation to pattern recognition. *IEEE Trans on Systems, Man, and Cybernetics SMC-15 No 5*: 608-619
- Paluš M (1993) Identifying and quantifying chaos by using information-theoretic functionals. In: Weigend AS, Gershenfeld NA (eds) *Time Series Prediction: Forecasting the Future and Understanding the Past*. Santa Fe Institute Studies in the Sciences of Complexity, Proc. Vol. XV. Addison-Wesley, Reading, Mass. pp 387-413
- Paluš M, Albrecht V, Dvořák I (1993) Information-theoretic test for nonlinearity in time series. *Phys Lett A* 175: 203-209
- Paluš M (1994a) Detection of nonlinearity and chaos in time series. Santa Fe Institute Working Paper 94-10-053 (<ftp://ftp.santafe.edu/pub/mp/eeg/a1-intro/>)
- Paluš M (1994b) Nonlinearity in normal human EEG: Cycles and randomness, not chaos. Santa Fe Institute Working Paper 94-10-054 (<ftp://ftp.santafe.edu/pub/mp/eeg/a2-res/>)

- Paluš M (1995) Testing for nonlinearity using redundancies: Quantitative and qualitative aspects. *Physica D* 80: 186-205
- Paluš M (submitted) Coarse-grained entropy rates for characterization of complex time series. Santa Fe Institute working paper 94-06-040; submitted to *Physica D* (<ftp://ftp.santafe.edu/pub/mp/eq/>)
- Pincus SM, Gladstone IM, Ehrenkranz RA (1991) A regularity statistic for medical data analysis. *J Clin Monit* 7: 335-345
- Pompe B (1993) Measuring statistical dependences in a time series. *J Stat Phys* 73(3/4): 587-610
- Press WH, Flannery BP, Teukolsky SA, Vetterling VT (1986) *Numerical Recipes: The Art of Scientific Computing*. Cambridge Univ Press, Cambridge
- Prichard D, Theiler J (1994) Generating surrogate data for time series with several simultaneously measured variables. *Phys Rev Lett* 73: 951-954
- Rapp PE, Bashore TR, Martinerie JM, Albano AM, Zimmerman ID, Mees AI (1989) Dynamics of brain electrical activity. *Brain Topography* 2(1/2): 99-118
- Theiler J, Eubank S, Longtin A, Galdrikian B, Farmer JD (1992) Testing for nonlinearity in time series: the method of surrogate data. *Physica D* 58: 77-94
- Theiler J, Rapp PE (1995) Re-examination of the evidence for low-dimensional nonlinear structure in the human electroencephalogram. *EEG Clin Neurophysiol*, in press
- Wackermann J, Lehmann D, Dvořák I, Michel CM (1993) Global dimensional complexity of multi-channel EEG indicates change of human brain functional state after a single dose of a nootropic drug. *EEG Clin Neurophysiol* 86: 193-198
- Weigend AS, Gershenfeld NA (eds) (1993) *Time Series Prediction: Forecasting the Future and Understanding the Past*. Santa Fe Institute Studies in the Sciences of Complexity, Proc. Vol. XV. Addison-Wesley, Reading, Mass.