

Two systems of point-free affine geometry

Giangiacomo Gerla ¹ Rafał Gruszczyński ²

¹IIASS
University of Salerno
Italy

²Department of Logic
Nicolaus Copernicus University in Toruń
Poland

TACL 2017

Outline

- 1 Inspirations and objectives
- 2 Half-plane structures
- 3 Oval structures

Outline

- 1 Inspirations and objectives
- 2 Half-plane structures
- 3 Oval structures

Inspirations and objectives

- 1 A. N. Whitehead and ovate class of regions
- 2 Aleksander Śniatycki and half-planes
- 3 affine geometry
- 4 follow geometrical intuitions
- 5 (regular) open convex subsets of \mathbb{R}^2 — «the litmus paper»

Inspirations and objectives

- 1 A. N. Whitehead and ovate class of regions
- 2 Aleksander Śniatycki and half-planes
- 3 affine geometry
- 4 follow geometrical intuitions
- 5 (regular) open convex subsets of \mathbb{R}^2 — «the litmus paper»

Inspirations and objectives

- 1 A. N. Whitehead and ovate class of regions
- 2 Aleksander Śniatycki and half-planes
- 3 affine geometry
- 4 follow geometrical intuitions
- 5 (regular) open convex subsets of \mathbb{R}^2 — «the litmus paper»

Inspirations and objectives

- 1 A. N. Whitehead and ovate class of regions
- 2 Aleksander Śniatycki and half-planes
- 3 affine geometry
- 4 follow geometrical intuitions
- 5 (regular) open convex subsets of \mathbb{R}^2 — «the litmus paper»

Inspirations and objectives

- 1 A. N. Whitehead and ovate class of regions
- 2 Aleksander Śniatycki and half-planes
- 3 affine geometry
- 4 follow geometrical intuitions
- 5 (regular) open convex subsets of \mathbb{R}^2 — «the litmus paper»

Inspirations and objectives

- 1 A. N. Whitehead and ovate class of regions
- 2 Aleksander Śniatycki and half-planes
- 3 affine geometry
- 4 follow geometrical intuitions
- 5 (regular) open convex subsets of \mathbb{R}^2 — «the litmus paper»

Outline

- 1 Inspirations and objectives
- 2 Half-plane structures
- 3 Oval structures

Basic notions of Śniatycki's approach

We begin with an examination of triples $\langle \mathbf{R}, \leq, \mathbf{H} \rangle$ in which:

- \mathbf{R} is a non-empty set whose elements are called **regions**,
- $\langle \mathbf{R}, \leq \rangle$ is a complete Boolean lattice,
- $\mathbf{H} \subseteq \mathbf{R}$ is a set whose elements are called **half-planes** (we assume that $\mathbf{1}$ and $\mathbf{0}$ are not half-planes).

Specific axioms for half-planes

$$h \in \mathbf{H} \longrightarrow -h \in \mathbf{H} \quad (\text{H1})$$

$$\begin{aligned} \forall_{x_1, x_2, x_3 \in \mathbf{R}} \big(\exists_{h \in \mathbf{H}} \forall_{i \in \{1, 2, 3\}} (x_i \cdot h \neq \mathbf{0} \wedge x_i \cdot -h \neq \mathbf{0}) \vee \\ \exists_{h_1, h_2, h_3 \in \mathbf{H}} (x_1 \leq h_1 \wedge x_2 \leq h_2 \wedge x_3 \leq h_3 \wedge \\ x_1 + x_2 \perp h_2 \wedge x_1 + x_3 \perp h_2 \wedge x_2 + x_3 \perp h_1) \big) \end{aligned} \quad (\text{H2})$$

Specific axioms for half-planes

$$h \in \mathbf{H} \longrightarrow -h \in \mathbf{H} \quad (\text{H1})$$

$$\begin{aligned} \forall_{x_1, x_2, x_3 \in \mathbf{R}} \big(\exists_{h \in \mathbf{H}} \forall_{i \in \{1, 2, 3\}} (x_i \cdot h \neq \mathbf{0} \wedge x_i \cdot -h \neq \mathbf{0}) \vee \\ \exists_{h_1, h_2, h_3 \in \mathbf{H}} (x_1 \leq h_1 \wedge x_2 \leq h_2 \wedge x_3 \leq h_3 \wedge \\ x_1 + x_2 \perp h_2 \wedge x_1 + x_3 \perp h_2 \wedge x_2 + x_3 \perp h_1) \big) \end{aligned} \quad (\text{H2})$$

Specific axioms for half-planes

$$\forall_{h_1, h_2, h_3 \in \mathbf{H}} (h_2 \leq h_1 \wedge h_3 \leq h_1 \longrightarrow h_2 \leq h_3 \vee h_3 \leq h_2) \quad (\text{H3})$$




Figure: In Beltrami-Klein model there are half-planes contained in a given one but incomparable in terms of \leq . In the picture above h_1 and h_2 are both parts of h , yet neither $h_1 \leq h_2$ nor $h_2 \leq h_1$. The purpose of (H3) is to ensure that parallelity of lines is a Euclidean relation.

Specific axioms for half-planes

$$\forall h_1, h_2, h_3 \in \mathbf{H} (h_2 \leq h_1 \wedge h_3 \leq h_1 \longrightarrow h_2 \leq h_3 \vee h_3 \leq h_2) \quad (\text{H3})$$




Figure: In Beltrami-Klein model there are half-planes contained in a given one but incomparable in terms of \leq . In the picture above h_1 and h_2 are both parts of h , yet neither $h_1 \leq h_2$ nor $h_2 \leq h_1$. The purpose of (H3) is to ensure that parallelity of lines is a Euclidean relation.

Lines and parallelity relation

Definition (of a line)

$L \in \mathcal{P}(\mathbf{H})$ is a **line** iff there is a half-plane h such that $L = \{h, -h\}$:

$$L \in \mathfrak{L} \stackrel{\text{df}}{\longleftrightarrow} \exists_{h \in \mathbf{H}} L = \{h, -h\}. \quad (\text{df } \mathfrak{L})$$

Definition (of parallelity relation)

$L_1, L_2 \in \mathfrak{L}$ are **parallel** iff there are half-planes $h \in L_1$ and $h' \in L_2$ which are disjoint:

$$L_1 \parallel L_2 \stackrel{\text{df}}{\longleftrightarrow} \exists_{h \in L_1} \exists_{h' \in L_2} h \perp h'. \quad (\text{df } \parallel)$$

In case L_1 and L_2 are not parallel we say they **intersect** and write: ' $L_1 \nparallel L_2$ '.

Angles and bowties. . .

Definition

- Given two intersecting lines L_1 and L_2 by **an angle** we understand a region x such that for $h_1 \in L_1$ and $h_2 \in L_2$ we have $x = h_1 \cdot h_2$:

x is an angle $\stackrel{\text{df}}{\longleftrightarrow} \exists_{L_1, L_2 \in \mathfrak{L}} (L_1 \nparallel L_2 \wedge \exists_{h_1 \in L_1} \exists_{h_2 \in L_2} x = h_1 \cdot h_2)$.

- An angle x is **opposite** to an angle y iff there are $h_1, h_2 \in \mathbf{H}$ such that $x = h_1 \cdot h_2$ and $y = -h_1 \cdot -h_2$.
- A **bowtie** is the sum of an angle and its opposite.

Notice that every pair $L_1 = \{h_1, -h_1\}$, $L_2 = \{h_2, -h_2\}$ of non-parallel lines determines exactly four pairwise disjoint angles: $h_1 \cdot h_2$, $h_1 \cdot -h_2$, $-h_1 \cdot h_2$ and $-h_1 \cdot -h_2$.

Angles and bowties. . .

Definition

- Given two intersecting lines L_1 and L_2 by **an angle** we understand a region x such that for $h_1 \in L_1$ and $h_2 \in L_2$ we have $x = h_1 \cdot h_2$:

x is an angle $\stackrel{\text{df}}{\longleftrightarrow} \exists_{L_1, L_2 \in \mathfrak{L}} (L_1 \nparallel L_2 \wedge \exists_{h_1 \in L_1} \exists_{h_2 \in L_2} x = h_1 \cdot h_2)$.

- An angle x is **opposite** to an angle y iff there are $h_1, h_2 \in \mathbf{H}$ such that $x = h_1 \cdot h_2$ and $y = -h_1 \cdot -h_2$.
- A **bowtie** is the sum of an angle and its opposite.

Notice that every pair $L_1 = \{h_1, -h_1\}$, $L_2 = \{h_2, -h_2\}$ of non-parallel lines determines exactly four pairwise disjoint angles: $h_1 \cdot h_2$, $h_1 \cdot -h_2$, $-h_1 \cdot h_2$ and $-h_1 \cdot -h_2$.

Angles and bowties. . .

Definition

- Given two intersecting lines L_1 and L_2 by **an angle** we understand a region x such that for $h_1 \in L_1$ and $h_2 \in L_2$ we have $x = h_1 \cdot h_2$:

x is an angle $\stackrel{\text{df}}{\longleftrightarrow} \exists_{L_1, L_2 \in \mathfrak{L}} (L_1 \nparallel L_2 \wedge \exists_{h_1 \in L_1} \exists_{h_2 \in L_2} x = h_1 \cdot h_2)$.

- An angle x is **opposite** to an angle y iff there are $h_1, h_2 \in \mathbf{H}$ such that $x = h_1 \cdot h_2$ and $y = -h_1 \cdot -h_2$.
- A **bowtie** is the sum of an angle and its opposite.

Notice that every pair $L_1 = \{h_1, -h_1\}$, $L_2 = \{h_2, -h_2\}$ of non-parallel lines determines exactly four pairwise disjoint angles: $h_1 \cdot h_2$, $h_1 \cdot -h_2$, $-h_1 \cdot h_2$ and $-h_1 \cdot -h_2$.

Angles and bowties. . .

Definition

- Given two intersecting lines L_1 and L_2 by **an angle** we understand a region x such that for $h_1 \in L_1$ and $h_2 \in L_2$ we have $x = h_1 \cdot h_2$:

x is an angle $\stackrel{\text{df}}{\longleftrightarrow} \exists_{L_1, L_2 \in \mathfrak{L}} (L_1 \nparallel L_2 \wedge \exists_{h_1 \in L_1} \exists_{h_2 \in L_2} x = h_1 \cdot h_2)$.

- An angle x is **opposite** to an angle y iff there are $h_1, h_2 \in \mathbf{H}$ such that $x = h_1 \cdot h_2$ and $y = -h_1 \cdot -h_2$.
- A **bowtie** is the sum of an angle and its opposite.

Notice that every pair $L_1 = \{h_1, -h_1\}$, $L_2 = \{h_2, -h_2\}$ of non-parallel lines determines exactly four pairwise disjoint angles: $h_1 \cdot h_2$, $h_1 \cdot -h_2$, $-h_1 \cdot h_2$ and $-h_1 \cdot -h_2$.

...and stripes

Definition

If $L_1 = \{h_1, -h_1\}$ and $L_2 = \{h_2, -h_2\}$ are parallel, yet distinct, lines and h_1 and h_2 are their disjoint sides, then $-h_1 \cdot -h_2$ is **stripe**.

Examples in the intended model




Figure: Fragments of a bowtie, a stripe and the complement of a stripe. These are all possible non-zero forms of the disjoint union of two distinct half-planes in the intended model. Any of the two shaded triangular areas of the bowtie is an angle.

Specific axioms for half-planes

$$\begin{aligned} h_1 \cdot h_2 \leq (h_3 \cdot h_4) + (-h_3 \cdot -h_4) \longrightarrow \\ h_3 = h_4 \vee h_1 \cdot h_2 \leq h_3 \cdot h_4 \vee h_1 \cdot h_2 \leq -h_3 \cdot -h_4 . \end{aligned} \quad (\text{H4})$$

Specific axioms for half-planes

$$h_1 \cdot h_2 \leq (h_3 \cdot h_4) + (-h_3 \cdot -h_4) \longrightarrow$$

$$h_3 = h_4 \vee h_1 \cdot h_2 \leq h_3 \cdot h_4 \vee h_1 \cdot h_2 \leq -h_3 \cdot -h_4. \quad (\text{H4})$$




Figure: A geometrical interpretation of (H4).

Specific axioms for half-planes

$$h_1 \cdot h_2 \leq (h_3 \cdot h_4) + (-h_3 \cdot -h_4) \longrightarrow$$

$$h_3 = h_4 \vee h_1 \cdot h_2 \leq h_3 \cdot h_4 \vee h_1 \cdot h_2 \leq -h_3 \cdot -h_4. \quad (\text{H4})$$




Figure: These two situations are excluded by the special case of (H4).

Points

Definition

Given lines L_1, \dots, L_k by a **net** determined by them we understand the following set:

$$(L_1 \dots L_k) := \{g_1 \cdot \dots \cdot g_k \mid \forall_{i \leq k} g_i \in L_i\}.$$

Lines L_1, \dots, L_k **split** a region x into m parts iff the set:

$$\{x \cdot a \mid a \in (L_1 \dots L_k) \wedge x \cdot a \neq \mathbf{0}\}$$

has exactly m elements.

Points

Definition

- If $L_1, \dots, L_k \in \mathfrak{L}$, an arbitrary element of the Cartesian product $L_1 \times \dots \times L_k$ will be called an *H-sequence*.
- An *H-sequence* $\langle h_1, \dots, h_k \rangle$ is *positive* iff $h_1 \cdot \dots \cdot h_k \neq \mathbf{0}$, otherwise it is *non-positive*.
- Two *H-sequences* $\langle g_1, \dots, g_k \rangle$ and $\langle g_1^*, \dots, g_k^* \rangle$ are *opposite* iff for all $i \leq n$, g_i^* is the complement of g_i .

Points

Definition

- If $L_1, \dots, L_k \in \mathfrak{L}$, an arbitrary element of the Cartesian product $L_1 \times \dots \times L_k$ will be called an *H-sequence*.
- An *H-sequence* $\langle h_1, \dots, h_k \rangle$ is *positive* iff $h_1 \cdot \dots \cdot h_k \neq \mathbf{0}$, otherwise it is *non-positive*.
- Two *H-sequences* $\langle g_1, \dots, g_k \rangle$ and $\langle g_1^*, \dots, g_k^* \rangle$ are *opposite* iff for all $i \leq n$, g_i^* is the complement of g_i .

Points

Definition

- If $L_1, \dots, L_k \in \mathfrak{L}$, an arbitrary element of the Cartesian product $L_1 \times \dots \times L_k$ will be called an *H-sequence*.
- An *H-sequence* $\langle h_1, \dots, h_k \rangle$ is *positive* iff $h_1 \cdot \dots \cdot h_k \neq \mathbf{0}$, otherwise it is *non-positive*.
- Two *H-sequences* $\langle g_1, \dots, g_k \rangle$ and $\langle g_1^*, \dots, g_k^* \rangle$ are *opposite* iff for all $i \leq n$, g_i^* is the complement of g_i .

Points

Definition

A **pseudopoint** is any net $(L_1 L_2)$ such that all its four H -sequences are positive, equivalently one could define a pseudopoint as an unordered pair of non-parallel lines.

For any pseudopoint $(L_1 L_2)$, the lines L_1 and L_2 will be called its **determinants**. In case we have two pseudopoints $(L_1 L_2)$ and $(L_1 L_3)$ we say that they **share a determinant** L_1 .

Points

Definition

Lines L_1 , L_2 and L_3 are **tied** iff $L_1 \times L_2 \times L_3$ contains two different non-positive and opposite H -sequences.

Definition

A pseudopoint $(L_1 L_2)$ **lies** on L_3 iff L_1 , L_2 and L_3 are tied.

Definition

Pseudopoints $(L_1 L_2)$ and $(L_3 L_4)$ are **collocated** (in symbols: $(L_1 L_2) \sim (L_3 L_4)$) iff $(L_1 L_2)$ lies on both L_3 and L_4 .

Points

Definition

Lines L_1 , L_2 and L_3 are **tied** iff $L_1 \times L_2 \times L_3$ contains two different non-positive and opposite H -sequences.

Definition

A pseudopoint $(L_1 L_2)$ **lies** on L_3 iff L_1 , L_2 and L_3 are tied.

Definition

Pseudopoints $(L_1 L_2)$ and $(L_3 L_4)$ are **collocated** (in symbols: $(L_1 L_2) \sim (L_3 L_4)$) iff $(L_1 L_2)$ lies on both L_3 and L_4 .

Points

Definition


Lines L_1 , L_2 and L_3 are **tyed** iff $L_1 \times L_2 \times L_3$ contains two different non-positive and opposite H -sequences.

Definition


A pseudopoint $(L_1 L_2)$ **lies** on L_3 iff L_1 , L_2 and L_3 are tied.

Definition


Pseudopoints $(L_1 L_2)$ and $(L_3 L_4)$ are **collocated** (in symbols: $(L_1 L_2) \sim (L_3 L_4)$) iff $(L_1 L_2)$ lies on both L_3 and L_4 .



h_1	h_2	h_3	P
h_1	h_2	$-h_3$	P
h_1	$-h_2$	h_3	N
h_1	$-h_2$	$-h_3$	P
$-h_1$	h_2	h_3	P
$-h_1$	h_2	$-h_3$	P
$-h_1$	$-h_2$	h_3	P
$-h_1$	$-h_2$	$-h_3$	P



h_1	h_2	h_3	P
h_1	h_2	$-h_3$	P
h_1	$-h_2$	h_3	N
h_1	$-h_2$	$-h_3$	N
$-h_1$	h_2	h_3	P
$-h_1$	h_2	$-h_3$	P
$-h_1$	$-h_2$	h_3	P
$-h_1$	$-h_2$	$-h_3$	P



h_1	h_2	h_3	P
h_1	h_2	$-h_3$	P
h_1	$-h_2$	h_3	N
h_1	$-h_2$	$-h_3$	P
$-h_1$	h_2	h_3	P
$-h_1$	h_2	$-h_3$	N
$-h_1$	$-h_2$	h_3	P
$-h_1$	$-h_2$	$-h_3$	P

Points

Definition

Collocation of pseudopoints is an equivalence relation, therefore points can be defined as its equivalence classes:

$$\Pi := \pi / \sim . \quad (\text{df } \Pi)$$

Incidence relation

Definition

$\alpha \in \Pi$ is **incident** with a line L iff there is a pseudopoint $(L_1 L_2) \in \alpha$ such that $(L_1 L_2)$ lies on L .

Betweenness relation

Definition

- $\alpha \in \Pi$ **lies in the half-plane** h iff there is $(L_1 L_2) \in \alpha$ such that for every $x \in (L_1 L_2)$, $x \cdot h \neq \mathbf{0}$.
- A line $L = \{h, -h\}$ **lies between** points α and β iff α lies in h and β lies in $-h$.

Definition

Points α, β and γ are **collinear** iff some three pseudopoints from, respectively, α, β and γ share a determinant L .

Betweenness relation

Definition

- $\alpha \in \Pi$ **lies in the half-plane** h iff there is $(L_1 L_2) \in \alpha$ such that for every $x \in (L_1 L_2)$, $x \cdot h \neq \mathbf{0}$.
- A line $L = \{h, -h\}$ **lies between** points α and β iff α lies in h and β lies in $-h$.

Definition

Points α, β and γ are **collinear** iff some three pseudopoints from, respectively, α, β and γ share a determinant L .

Betweenness relation

Definition

- $\alpha \in \Pi$ **lies in the half-plane** h iff there is $(L_1 L_2) \in \alpha$ such that for every $x \in (L_1 L_2)$, $x \cdot h \neq \mathbf{0}$.
- A line $L = \{h, -h\}$ **lies between** points α and β iff α lies in h and β lies in $-h$.

Definition

Points α, β and γ are **collinear** iff some three pseudopoints from, respectively, α, β and γ share a determinant L .

Betweenness relation

Definition

A point γ is **between** points α and β iff:

- α, β and γ are collinear and
- γ is incident with a line L which lies between α and β .

Śniatycki's Theorem

Theorem

Consider an H -structure:

$$\langle \mathbf{R}, \leq, \mathbf{H} \rangle .$$

Individual notions of point and line and relational notions of incidence and betweenness are definable in such a way that the corresponding structure $\langle \Pi, \mathcal{L}, \epsilon, \mathbf{B} \rangle$ satisfies all axioms of a system of geometry of betweenness and incidence.

Outline

- 1 Inspirations and objectives
- 2 Half-plane structures
- 3 Oval structures

Basic notions

We now turn our attentions to structures $\langle \mathbf{R}, \leq, \mathbf{O} \rangle$ such that:

- elements of \mathbf{R} are called **regions**,
- $\leq \subseteq \mathbf{R}^2$ is **part of** relation,
- $\mathbf{O} \subseteq \mathbf{R}$ and its elements are called **ovals**.

First axioms

$\langle \mathbf{R}, \leq \rangle$ is a complete atomless Boolean lattice. (00)

\mathbf{O} is an algebraic closure system in $\langle \mathbf{R}, \leq \rangle$ containing $\mathbf{0}$. (01)

\mathbf{O}^+ is dense in $\langle \mathbf{R}, \leq \rangle$. (02)

The hull operator

Definition

$\text{hull}: \mathbf{R} \longrightarrow \mathbf{R}$ is the operation given by:

$$\text{hull}(x) := \bigwedge \{a \in \mathbf{O} \mid x \leq a\}. \quad (\text{df hull})$$

For $x \in \mathbf{R}$ the object $\text{hull}(x)$ will be called **the oval generated by x** .

Lines in the oval setting

Definition

By **a line** we understand a two element set $L = \{a, b\}$ of disjoint ovals, such that for any set of disjoint ovals $\{c, d\}$ with $a \leq c$ and $b \leq d$ it is the case that $a = c$ and $b = d$:

$$X \in \mathfrak{L} \stackrel{\text{df}}{\iff} \exists_{a,b \in \mathbf{O}^+} (a \perp b \wedge X = \{a, b\} \wedge \forall_{c,d \in \mathbf{O}^+} (c \perp d \wedge a \leq c \wedge b \leq d \longrightarrow a = c \wedge b = d)). \quad (\text{df } \mathfrak{L})$$

For a line $L = \{a, b\}$ the elements of L will be called **the sides of L** .

Lines in the oval setting

Definition

Two lines $L_1 = \{a, b\}$ and $L_2 = \{c, d\}$ are **parallel** iff there is a side of L_1 which is disjoint from a side of L_2 :

$$L_1 \parallel L_2 \stackrel{\text{df}}{\iff} \exists a \in L_1 \exists b \in L_2 \ a \perp b. \quad (\text{df } \parallel)$$

In case L_1 is not parallel to L_2 we say that L_1 and L_2 *intersect* and write ' $L_1 \nparallel L_2$ '.

Half-planes in the oval setting

Definition

A region x is a **half-plane** iff $x, -x \in \mathbf{O}^+$; the set of all half-planes will be denoted by ' \mathbf{H} ':

$$x \in \mathbf{H} \stackrel{\text{df}}{\iff} \{x, -x\} \subseteq \mathbf{O}^+ . \quad (\text{df } \mathbf{H})$$

Half-planes and lines in oval setting

Definition

Let B_1, \dots, B_n be non-empty spheres in \mathbb{R}^2 such that for $1 \leq i \neq j \leq n$: $\text{Cl } B_i \cap \text{Cl } B_j = \emptyset$. Consider the subspace \mathcal{B}_n of \mathbb{R}^2 induced by $B_1 \cup \dots \cup B_n$. Put:

- $\text{r}\mathcal{B}_n := \{x \mid x \text{ is a regular open element of } \mathcal{B}_n\}$
- $\mathbf{O} := \{a \in \text{r}\mathcal{B}_n \mid a = \bigcup_{1 \leq i \leq n} B_i \vee \exists_{1 \leq i \leq n} \exists_{b \in \text{Conv}} a = B_i \cap b\}$

We will call $\mathbb{B}_n := \langle \text{r}\mathcal{B}_n, \subseteq, \mathbf{O} \rangle$ the *n -sphere structure*.

Lines and half-planes in the oval setting




Figure: The structure \mathbb{B}_3 .

Lines and half-planes in the oval setting




Figure: The structure \mathbb{B}_3 .

Fact

For every $n \in \mathbb{N}$, \mathbb{B}_n is a complete Boolean lattice and the axioms (01) and (02) are satisfied in \mathbb{B}_n .

Lines and half-planes in the oval setting




Figure: The structure \mathbb{B}_3 .

Fact

For every $n \in \mathbb{N}$, the set of lines of \mathbb{B}_n contains sets $\{B_i \cap h, B_i \cap -h\}$, where h is a half-plane in the prototypical structure \mathbb{R}^2 and both $B_i \cap h$ and $B_i \cap -h$ are non-empty. Two lines contained in different balls are always parallel.

Lines and half-planes in the oval setting




Figure: The structure \mathbb{B}_1 .

Fact

In \mathbb{B}_1 the set of lines is equal to the set of all unordered pairs of the form $\{B_1 \cap h, B_1 \cap -h\}$. The sides of a line in \mathbb{B}_1 are half-planes in this structure.

Lines and half-planes in the oval setting




Figure: The structure \mathbb{B}_2 .

Fact

B_1 and B_2 are the only half-planes of \mathbb{B}_2 and thus $\{B_1, B_2\}$ is the only line of \mathbb{B}_2 whose sides are half-planes. This line is parallel to every other line. In general, in \mathbb{B}_n for $n \geq 2$ any pair $\{B_i, B_j\}$ with $i \neq j$ is a line parallel to every line in \mathbb{B}_n .

Lines and half-planes in the oval setting




Figure: The structure \mathbb{B}_3 .

Fact

There are no half-planes in \mathbb{B}_n for $n \geq 3$, and thus there are no lines whose sides are half-planes.

Specific axioms

Definition

A **finite partition** of the universe **1** is a set $\{x_1, \dots, x_n\} \subseteq \mathbf{R}$ whose elements are pairwise disjoint and such that $\bigvee \{x_1, \dots, x_n\} = \mathbf{1}$. For a partition $P = \{x_1, \dots, x_n\}$ and $x \in \mathbf{R}$ by *the partition of x induced by P* we understand the following set:

$$\{x \cdot x_i \mid 1 \leq i \leq n \wedge x \cdot x_i \neq \mathbf{0}\}.$$

The sides of a line form a partition of **1**; equivalently: the sides of a line are half-planes.

(03)

Specific axioms

Definition

A **finite partition** of the universe **1** is a set $\{x_1, \dots, x_n\} \subseteq \mathbf{R}$ whose elements are pairwise disjoint and such that $\bigvee \{x_1, \dots, x_n\} = \mathbf{1}$. For a partition $P = \{x_1, \dots, x_n\}$ and $x \in \mathbf{R}$ by *the partition of x induced by P* we understand the following set:

$$\{x \cdot x_i \mid 1 \leq i \leq n \wedge x \cdot x_i \neq \mathbf{0}\}.$$

The sides of a line form a partition of **1**; equivalently: the sides of a line are half-planes. (03)

Specific axioms

For any $a, b, c \in \mathbf{O}$ which are not aligned there is a line (04)
which separates a from $\text{hull}(b + c)$.

Specific axioms

If distinct lines L_1 and L_2 both cross an oval a , then they split a in at least three parts. (05)



Figure: L_1 and L_2 split the oval into 3 parts, while L_3 and L_4 split it into 4 parts.

Specific axioms

If distinct lines L_1 and L_2 both cross an oval a , then they split a in at least three parts. (05)




Figure: L_1 and L_2 split the oval into 3 parts, while L_3 and L_4 split it into 4 parts.

Specific axioms

No half-plane is part of any stripe and any angle. (06)

The purpose of (06) is to prove that parallelity of lines is transitive.




Figure: In Beltrami-Klein model: h is a part of the angle $h_2 \cdot -h_1$.

Specific axioms

No half-plane is part of any stripe and any angle. (06)

The purpose of (06) is to prove that parallelity of lines is transitive.




Figure: In Beltrami-Klein model: h is a part of the angle $h_2 \cdot -h_1$.

O-structures

Definition

A triple $\langle \mathbf{R}, \leq, \mathbf{O} \rangle$ is an **O-structure** iff $\langle \mathbf{R}, \leq, \mathbf{O} \rangle$ satisfies axioms (00)–(06).

Main theorems

Theorem

Let $\mathfrak{O} = \langle \mathbf{R}, \leq, \mathbf{O} \rangle$ be an O -structure and $\mathfrak{O}' := \langle \mathbf{R}, \leq, \mathbf{O}, \mathbf{H} \rangle$ be the structure obtained from \mathfrak{O} by defining \mathbf{H} as the set of all ovals whose complements are ovals. Then \mathfrak{O}' satisfies all axioms for H -structures.

Theorem

If \mathfrak{O}' is the extension of an O -structure \mathfrak{O} , then individual notions of point and line and relational notions of incidence and betweenness are definable from the operations and notions of \mathfrak{O}' in such a way that all the axioms of a system of affine geometry are satisfied by the corresponding structure $\langle \mathbf{P}, \mathfrak{L}, \epsilon, \mathbf{B} \rangle$.

Main theorems

Theorem

Let $\mathfrak{O} = \langle \mathbf{R}, \leq, \mathbf{O} \rangle$ be an O -structure and $\mathfrak{O}' := \langle \mathbf{R}, \leq, \mathbf{O}, \mathbf{H} \rangle$ be the structure obtained from \mathfrak{O} by defining \mathbf{H} as the set of all ovals whose complements are ovals. Then \mathfrak{O}' satisfies all axioms for H -structures.

Theorem

If \mathfrak{O}' is the extension of an O -structure \mathfrak{O} , then individual notions of point and line and relational notions of incidence and betweenness are definable from the operations and notions of \mathfrak{O}' in such a way that all the axioms of a system of affine geometry are satisfied by the corresponding structure $\langle \mathbf{P}, \mathfrak{L}, \epsilon, \mathbf{B} \rangle$.

A proof of (H4)

We prove more general statement according to which for any $a \in \mathbf{O}$:

$$a \leq (h_3 \cdot h_4) + (-h_3 \cdot -h_4) \longrightarrow h_3 = h_4 \vee a \leq h_3 \cdot h_4 \vee a \leq -h_3 \cdot -h_4 ,$$

and use the fact that for any half planes h_1 and h_2 , $h_1 \cdot h_2 \in \mathbf{O}$.

The case in which $a = \mathbf{0}$ is trivial. In case $h_3 = -h_4$ we have that:

$$(h_3 \cdot h_4) + (-h_3 \cdot -h_4) = (-h_4 \cdot h_4) + (h_4 \cdot -h_4) = \mathbf{0} .$$

Thus we assume that (a) $h_3 \neq -h_4$. Let:

$$a \leq (h_3 \cdot h_4) + (-h_3 \cdot -h_4) \quad (\bullet)$$

and (b) $h_3 \neq h_4$. At the same time assume towards contradiction that:

$$a \not\leq h_3 \cdot h_4 \quad \text{and} \quad a \not\leq -h_3 \cdot -h_4 . \quad (\ddagger)$$

A proof of (H4)

By (a) and (b) lines $L_3 = \{h_3, -h_3\}$ and $L_4 = \{h_4, -h_4\}$ are distinct. From (\bullet) and (\ddagger) we get that $a \cdot h_3 \cdot h_4 \neq \mathbf{0} \neq a \cdot -h_3 \cdot -h_4$, so both L_3 and L_4 cross a and according to axiom (O5) they split a into at least three parts. Yet (\bullet) entails that:

$$a \cdot -h_3 \cdot h_4 \leq ((h_3 \cdot h_4) + (-h_3 \cdot -h_4)) \cdot -h_3 \cdot h_4 = \mathbf{0}$$

and

$$a \cdot h_3 \cdot -h_4 \leq ((h_3 \cdot h_4) + (-h_3 \cdot -h_4)) \cdot h_3 \cdot -h_4 = \mathbf{0}$$

and in consequence the set:

$$\{a \cdot x \mid x \in (L_3 L_4) \wedge a \cdot x \neq \mathbf{0}\}$$

has exactly two elements, a contradiction. □

Bibliography

Giorgio Gerla and Rafał Gruszczyński, **Point-free geometry, ovals and half-planes**, *Review of Symbolic Logic*, Volume 10, Issue 2 (2017), pp. 237–258

Aleksander Śniatycki, **An axiomatics of non-Desarguean geometry based on the half-plane as the primitive notion**, *Dissertationes Mathematicae*, no. LIX, PWN, Warszawa, 1968

Support

Research supported by National Science Center, Poland, grant
Applications of mereology in systems of point-free geometry, no.
2014/13/B/HS1/00766.

The End